USTC-ICTS-16-04

Wess-Zumino Model on Bosonic-Fermionic Noncommutative Superspace

Xu-Dong Wang[1]

Interdisciplinary Center for Theoretical Study
University of Science and Technology of China, Hefei, Anhui 230026, People's Republic of China

arXiv:1603.06302v1 [hep-th] 21 Mar 2016

Abstract

In our previous paper we construct a renormalizable Wess-Zumino action on BFNC superspace at the second order approximation of noncommutative parameters. The action contains about 200 terms which are necessary for renormalization. By removing chiral covariant derivatives and chiral coordinates we found that the BFNC Wess-Zumino action can be transformed to a simpler form which have manifest 1/2 supersymmetry. Based on this discovery, we can extend the BFNC Wess-Zumino action to the all order of noncommutative parameters. At first we introduce global symmetries, then obtain divergent operators in the effective action by using dimensional analysis, the next step is to construct all possible BFNC parameters, at the end we combine the BFNC parameters with the divergent operators. We present the explicit action up to the fourth order of noncommutative parameters. Because the action contain all possible divergent operators, it is renormalizable to all order in perturbative theory.

PACS Number(s): 12.60.Jv, 11.30.Pb, 11.10.Nx, 11.10.Gh

Keywords: Wess-Zumino Model, Bosonic-Fermionic Noncommutative Superspace, Renormalization

[1]E-mail address: xudwang@ustc.edu.cn

1 Introduction

There are many studies on noncommutative field theory in recent years [1, 2]. The introduction of supersymmetry on noncommutative spacetime is also very interesting. We can classify the noncommutative superspace by three kind. The first kind is Bosonic noncommutative superspace in which the algebraic relation is $\left[x^k,\ x^l\right] = i\ \theta^{kl}$. It can be obtained from string theory [3]. Ref. [4] constructed Wess-Zumino and Super Yang-Mills model on this superspace. The renormalization property of these models were studied in Ref. [5]. They proved that the Wess-Zumino model is renormalizable.

The second kind is non-anticommutative (NAC) superspace $\left\{\theta^\alpha,\ \theta^\beta\right\} = C^{\alpha\beta}$. It can also be obtained from string theory [6, 7]. In Ref. [8] the author defined Wess-Zumino and Super Yang-Mills model on this superspace. There are many work studying these models, for example Ref. [9] calculate the 1PI effective action for NAC Wess-Zumino model, and found a new term $(-g^2/\epsilon)\int d^8z\ \ (m^*)^2\, U\left(D^2\Phi\right)^2$, where $U = \theta^2\ \bar{\theta}^2\ C^2$ in the effective action. This term make the deformed action not renormalizable. By introducing two global $U(1)$ symmetries, Ref. [10, 11] prove that if one add this term to the deformed action, the new action is renormalizable to all order in perturbative theory.

The third kind is Bosonic-Fermionic noncommutative (BFNC) superspace, the algebraic relation is $\left[x^k,\ \theta^\alpha\right] = i\ \Lambda^{k\alpha}$. In Ref. [12] the author found the origin of this structure in string theory. To study the field theory on BFNC superspace, in Ref. [13, 14], we constructed a deformed Wess-Zumino model. By calculating the 1PI effective action, we obtain three effective action $\Gamma\left(\Lambda^2\right) = a_0\,\mathcal{S}_\Lambda\left(\Lambda^2\right) + a_1\,\Gamma_{\rm 1st}\left(\Lambda^2\right) + a_2\,\Gamma_{\rm 2nd}\left(\Lambda^2\right) + a_3\,\Gamma_{\rm 3rd}\left(\Lambda^2\right)$ and classified them by 1/2 supersymmetry invariant subset $\Gamma\left(\Lambda^2\right) = \sum_{i=1}^{74} f_i$. By using these invariant subset, we obtain more general 1/2 supersymmetry invariant action $\mathcal{S}_{(3)} = \mathcal{S}_{\rm WZ} + \int d^8z\left(\sum_{i=1}^{74} B_i\right)$ and prove that it is one-loop renormalizable. To extend the renormalizability to all order in perturbative theory, we introduce two global $U(1)$ symmetries and obtain a modified action $\mathcal{S}'_{(3)} = \mathcal{S}'_{\rm WZ} + \int d^8z\left(\sum_{i=1}^{74} c_i\ B_i\right)$, after adding the other divergent terms obtained by using dimensional analysis we obtain a BFNC Wess-Zumino model $\mathcal{S}'_{(3)} + \int d^8z\left(\sum_{i=75}^{89} c_i\ B_i\right)$ at the second order approximation of noncommutative parameters. Because the action contain all of the divergent operators, it is renormalizable to all order in perturbative theory.

In our previous studies [13, 14] the action contains only second order of noncommutative parameters. The associate property of BFNC star product need higher order of noncommutative parameters. Even at second order of noncommutative parameters, the BFNC Wess-Zumino model contain about 200 terms, this make the extension to higher oder approximation of

noncommutative parameters very difficult. Inspired by recent studies on super Yang-Mills theory on NAC superspace [15] and BFNC superspace [16], we found that by removing chiral covariant derivatives D^2 and chiral coordinates θ^2, the BFNC Wess-Zumino action can be transformed to more simpler form. With this discovery, we can extend the BFNC Wess-Zumino action to the all order of noncommutative parameters. To illustrate the idea, we will present the explicit action up to the fourth order of noncommutative parameters in this paper.

The organization of this paper is as follows, in section 2 we determine the divergent operators by using dimensional analysis method up to the fourth order of noncommutative parameters, in section 3 we construct all related BFNC parameters, in section 4 we construct 1/2 supersymmetric invariant action by combining the BFNC parameters with the divergent operators, we give our conclusion and outlook in section 5.

2 Divergent operators

By using two global $U(1)$ symmetries, we can determine the divergent operators in the effective action. We list them as follows, at the order of Λ^2,

- $N_\Phi = N_{D^2},\ N_{\bar{D}} = 0,\ N_{\partial\partial} = 0.$

$$\begin{aligned}&\Phi^+, \quad \Phi^+\Phi^+, \quad \Phi^+\Phi^+\Phi^+, \quad \Phi^+\Phi^+\Phi^+\Phi^+, \quad \Phi^+\Phi^+\Phi^+\Phi^+\Phi^+,\\ &D^2\Phi, \quad D^2\Phi\Phi^+, \quad D^2D^2\Phi\Phi, \quad D^2\Phi\Phi^+\Phi^+, \quad D^2D^2\Phi\Phi\Phi^+, \quad D^2\Phi\Phi^+\Phi^+\Phi^+.\end{aligned} \tag{1}$$

- $N_\Phi = N_{D^2},\ N_{\bar{D}} = 0,\ N_{\partial\partial} = 1.$

$$\begin{aligned}&\partial\partial\Phi^+\Phi^+, \quad D^2\partial\partial\Phi\Phi^+, \quad D^2D^2\partial\partial\Phi\Phi, \quad \partial\partial\Phi^+\Phi^+\Phi^+, \quad D^2\partial\partial\Phi\Phi^+\Phi^+,\\ &\partial\partial\Phi^+\Phi^+\Phi^+\Phi^+.\end{aligned} \tag{2}$$

- $N_\Phi = N_{D^2},\ N_{\bar{D}} = 0,\ N_{\partial\partial} = 2.$

$$\partial\partial\partial\partial\Phi^+\Phi^+, \quad D^2\partial\partial\partial\partial\Phi\Phi^+, \quad \partial\partial\partial\partial\Phi^+\Phi^+\Phi^+. \tag{3}$$

- $N_\Phi = N_{D^2},\ N_{\bar{D}} = 2,\ N_{\partial\partial} = 0.$

$$\begin{aligned}&\bar{D}^2\Phi^+, \quad \bar{D}^2\Phi^+\Phi^+, \quad \bar{D}^2\Phi^+\Phi^+\Phi^+, \quad \bar{D}^2\Phi^+\Phi^+\Phi^+\Phi^+, \quad \bar{D}^2\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+,\\ &\bar{D}^2\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+.\end{aligned} \tag{4}$$

- $N_\Phi > N_{D^2},\ N_{\bar{D}} = 0,\ N_{\partial\partial} = 0.$

$$D^2\Phi\Phi, \quad D^2\Phi\Phi\Phi, \quad D^2D^2\Phi\Phi\Phi, \quad D^2\Phi\Phi\Phi^+,$$

$$D^2D^2\Phi\Phi\Phi\Phi, \quad D^2\Phi\Phi\Phi\Phi^+, \quad D^2D^2\Phi\Phi\Phi\Phi^+, \quad D^2\Phi\Phi\Phi^+\Phi^+, \quad D^2\Phi\Phi\Phi^+\Phi^+\Phi^+,$$
$$D^2\Phi\Phi\Phi\Phi^+\Phi^+, \quad D^2D^2\Phi\Phi\Phi\Phi\Phi^+, \quad D^2\Phi\Phi\Phi\Phi^+\Phi^+\Phi^+. \tag{5}$$

- The other operators,

$$D^2\partial\partial\Phi\Phi, \quad D^2D^2\partial\partial\Phi\Phi\Phi,$$
$$D^2\overline{D}^2\Phi\Phi^+, \quad \partial D\bar{D}\Phi\Phi^+, \quad \partial\partial\Phi\Phi^+,$$
$$D^2\overline{D}^2\Phi\Phi^+\Phi^+, \quad \partial D\bar{D}\Phi\Phi^+\Phi^+, \quad \partial\partial\Phi\Phi^+\Phi^+,$$
$$D^2D^2\overline{D}^2\Phi\Phi\Phi^+, \quad D^2\partial D\bar{D}\Phi\Phi\Phi^+, \quad D^2\partial\partial\Phi\Phi\Phi^+,$$
$$D^2\overline{D}^2\Phi\Phi^+\Phi^+\Phi^+, \quad \partial D\bar{D}\Phi\Phi^+\Phi^+\Phi^+, \quad \partial\partial\Phi\Phi^+\Phi^+\Phi^+,$$
$$D^2D^2\overline{D}^2\Phi\Phi\Phi^+\Phi^+, \quad D^2\partial D\bar{D}\Phi\Phi\Phi^+\Phi^+, \quad D^2\partial\partial\Phi\Phi\Phi^+\Phi^+,$$
$$D^2\overline{D}^2\Phi\Phi^+\Phi^+\Phi^+\Phi^+, \quad \partial D\bar{D}\Phi\Phi^+\Phi^+\Phi^+\Phi^+, \quad \partial\partial\Phi\Phi^+\Phi^+\Phi^+\Phi^+. \tag{6}$$

At the order of Λ^4,

- $N_\Phi = N_{D^2},\ N_{\bar{D}} = 0,\ N_{\partial\partial} = 0.$

$$D^2D^2D^2\Phi\Phi\Phi,$$
$$D^2D^2\Phi\Phi\Phi^+\Phi^+, \quad D^2D^2D^2\Phi\Phi\Phi\Phi^+, \quad D^2D^2D^2D^2\Phi\Phi\Phi\Phi,$$
$$D^2\Phi\Phi^+\Phi^+\Phi^+\Phi^+, \quad D^2D^2\Phi\Phi\Phi^+\Phi^+\Phi^+, \quad D^2D^2D^2\Phi\Phi\Phi\Phi^+\Phi^+,$$
$$D^2\Phi\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+, \quad D^2D^2\Phi\Phi\Phi^+\Phi^+\Phi^+\Phi^+,$$
$$D^2\Phi\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+. \tag{7}$$

- $N_\Phi = N_{D^2},\ N_{\bar{D}} = 0,\ N_{\partial\partial} = 1.$

$$D^2D^2\partial\partial\Phi\Phi\Phi\Phi^+, \quad D^2D^2D^2\partial\partial\Phi\Phi\Phi,$$
$$D^2\partial\partial\Phi\Phi^+\Phi^+\Phi^+, \quad D^2D^2\partial\partial\Phi\Phi\Phi^+\Phi^+, \quad D^2D^2D^2\partial\partial\Phi\Phi\Phi\Phi^+,$$
$$\partial\partial\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+, \quad D^2\partial\partial\Phi\Phi^+\Phi^+\Phi^+\Phi^+, \quad D^2D^2\partial\partial\Phi\Phi\Phi^+\Phi^+\Phi^+,$$
$$\partial\partial\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+, \quad D^2\partial\partial\Phi\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+,$$
$$\partial\partial\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+. \tag{8}$$

- $N_\Phi = N_{D^2},\ N_{\bar{D}} = 0,\ N_{\partial\partial} = 2.$

$$D^2D^2\partial\partial\partial\partial\Phi\Phi,$$
$$D^2\partial\partial\partial\partial\Phi\Phi^+\Phi^+, \quad D^2D^2\partial\partial\partial\partial\Phi\Phi\Phi^+, \quad D^2D^2D^2\partial\partial\partial\partial\Phi\Phi\Phi,$$
$$\partial\partial\partial\partial\Phi^+\Phi^+\Phi^+\Phi^+, \quad D^2\partial\partial\partial\partial\Phi\Phi^+\Phi^+\Phi^+, \quad D^2D^2\partial\partial\partial\partial\Phi\Phi\Phi^+\Phi^+,$$
$$\partial\partial\partial\partial\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+, \quad D^2\partial\partial\partial\partial\Phi\Phi^+\Phi^+\Phi^+\Phi^+,$$
$$\partial\partial\partial\partial\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+. \tag{9}$$

- $N_\Phi > N_{D^2},\ N_{\bar{D}} = 0,\ N_{\partial\partial} = 0.$

$$
\begin{aligned}
&D^2D^2D^2\Phi\Phi\Phi\Phi,\\
&D^2D^2\Phi\Phi\Phi\Phi^+\Phi^+,\quad D^2D^2D^2\Phi\Phi\Phi\Phi\Phi,\quad D^2D^2D^2\Phi\Phi\Phi\Phi\Phi^+,\quad D^2D^2D^2D^2\Phi\Phi\Phi\Phi\Phi,\\
&D^2\Phi\Phi\Phi^+\Phi^+\Phi^+\Phi^+,\quad D^2D^2\Phi\Phi\Phi\Phi^+\Phi^+\Phi^+,\quad D^2D^2\Phi\Phi\Phi\Phi\Phi^+\Phi^+,\\
&D^2D^2D^2\Phi\Phi\Phi\Phi\Phi^+\Phi^+,\quad D^2D^2D^2\Phi\Phi\Phi\Phi\Phi\Phi^+,\quad D^2D^2D^2D^2\Phi\Phi\Phi\Phi\Phi\Phi,\\
&D^2\Phi\Phi\Phi^+\Phi^+\Phi^+\Phi^+,\quad D^2\Phi\Phi\Phi\Phi^+\Phi^+\Phi^+\Phi^+,\quad D^2D^2\Phi\Phi\Phi\Phi^+\Phi^+\Phi^+\Phi^+,\\
&D^2D^2\Phi\Phi\Phi\Phi\Phi^+\Phi^+\Phi^+,\quad D^2D^2D^2\Phi\Phi\Phi\Phi\Phi\Phi\Phi^+\Phi^+,\\
&D^2\Phi\Phi\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+,\quad D^2\Phi\Phi\Phi\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+,\quad D^2D^2\Phi\Phi\Phi\Phi\Phi^+\Phi^+\Phi^+\Phi^+,\\
&D^2\Phi\Phi\Phi\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+.
\end{aligned}
\tag{10}
$$

- The other operators,

$$
\begin{aligned}
&D^2D^2D^2\partial\partial\Phi\Phi\Phi\Phi,\\
&D^2D^2D^2\overline{D}^2\Phi\Phi\Phi\Phi^+,\quad D^2D^2\partial D\overline{D}\Phi\Phi\Phi\Phi^+, D^2D^2\partial\partial\Phi\Phi\Phi\Phi^+,\\
&D^2D^2\overline{D}^2\Phi\Phi\Phi^+\Phi^+\Phi^+,\quad D^2\partial D\overline{D}\Phi\Phi\Phi^+\Phi^+\Phi^+,\quad D^2\partial\partial\Phi\Phi\Phi^+\Phi^+\Phi^+,\\
&D^2D^2D^2\overline{D}^2\Phi\Phi\Phi\Phi^+\Phi^+,\quad D^2D^2\partial D\overline{D}\Phi\Phi\Phi\Phi^+\Phi^+,\quad D^2D^2\partial\partial\Phi\Phi\Phi\Phi^+\Phi^+,\\
&D^2D^2D^2D^2\overline{D}^2\Phi\Phi\Phi\Phi\Phi^+,\quad D^2D^2D^2\partial D\overline{D}\Phi\Phi\Phi\Phi\Phi^+,\quad D^2D^2D^2\partial\partial\Phi\Phi\Phi\Phi\Phi^+,\\
&D^2\overline{D}^2\Phi\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+,\quad \partial D\overline{D}\Phi\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+,\quad \partial\partial\Phi\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+,\\
&D^2D^2\overline{D}^2\Phi\Phi\Phi^+\Phi^+\Phi^+\Phi^+,\quad D^2\partial D\overline{D}\Phi\Phi\Phi^+\Phi^+\Phi^+\Phi^+,\quad D^2\partial\partial\Phi\Phi\Phi^+\Phi^+\Phi^+\Phi^+,\\
&D^2D^2D^2\overline{D}^2\Phi\Phi\Phi\Phi^+\Phi^+\Phi^+,\quad D^2D^2\partial D\overline{D}\Phi\Phi\Phi\Phi^+\Phi^+\Phi^+,\quad D^2D^2\partial\partial\Phi\Phi\Phi\Phi^+\Phi^+\Phi^+,\\
&D^2\overline{D}^2\Phi\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+,\quad \partial D\overline{D}\Phi\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+,\quad \partial\partial\Phi\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+,\\
&D^2D^2\overline{D}^2\Phi\Phi\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+,\quad D^2\partial D\overline{D}\Phi\Phi\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+,\quad D^2\partial\partial\Phi\Phi\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+,\\
&D^2\overline{D}^2\Phi\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+,\quad \partial D\overline{D}\Phi\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+,\quad \partial\partial\Phi\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+\Phi^+.
\end{aligned}
\tag{11}
$$

3 BFNC parameters

In order to combine with possible divergent operators, we need BFNC parameters with the following indices,

$$k\ l,\quad \alpha\ \beta,\quad \dot{\alpha}\ \dot{\beta},\quad k\ \dot{\alpha}\ \beta,\quad k\ l\ \alpha\ \beta,\quad k\ l\ n\ o, \tag{12}$$

they can only be constructed by using the following elements,

$$\epsilon_{\alpha\beta},\quad \epsilon^{\alpha\beta},\quad \epsilon_{\dot{\alpha}\dot{\beta}},\quad \epsilon^{\dot{\alpha}\dot{\beta}},\quad \eta_{kl},\quad \eta^{kl},$$

$$\epsilon^{klmn}, \quad (\sigma^{kl})^{\alpha\beta}, \quad (\bar{\sigma}^{k})^{\dot{\alpha}\beta}, \quad \Lambda^{k}{}_{\alpha}. \tag{13}$$

By using the algebraic relations of covariant derivatives D and $\bar{D}$, we found some BFNC parameters have certain symmetries. For example, if the BFNC parameter contains only α β and $\dot{\alpha}$ $\dot{\beta}$, it must be antisymmetric with respect to these indices. On the other hand, we can obtain some BFNC parameters from others. For example, if we combine parameters with indices k l α β and parameters with indices $\epsilon_{\alpha\beta}$ or η_{kl}, we can obtain parameters with indices k l or α β. The parameters with indices k l can also be obtained from parameters with indices k l n o. Because the indices $\dot{\alpha}$ β can be projected on $\left(\bar{\sigma}^{k}\right)^{\dot{\alpha}\beta}$, if we find all of the parameters with indices k l, then combine them with $\left(\bar{\sigma}^{k}\right)^{\dot{\alpha}\beta}$ by using η_{kl}, we will obtain all parameters with indices k $\dot{\alpha}$ β Because the indices k l α β can be projected on $\epsilon^{\alpha\beta}$ and $\left(\sigma^{kl}\right)^{\alpha\beta}$, to construct parameters with indices k l α β, we can combine $\epsilon^{\alpha\beta}$ and parameters with indices k l, or combine $\left(\sigma^{kl}\right)^{\alpha\beta}$ and parameters with indices k l or k l n o. We do not need to consider parameters with more Bosonic indices, because combining $\left(\sigma^{kl}\right)^{\alpha\beta}$ and these parameters equivalent to combining $\left(\sigma^{kl}\right)^{\alpha\beta}$ and parameters with indices k l or k l n o.

Based on above consideration, the task to construct parameters with indices in Eq. (12) is equivalent to constructing all parameters with indices k l n o. We give the details in Appendix. D and only list the BFNC parameters at order Λ^0, Λ^2 and Λ^4.

- At order Λ^0, parameters with indices kl and $klno$,

$$\eta^{kl}; \qquad \eta^{kl}\ \eta^{no}, \quad \epsilon^{klno}. \tag{14}$$

- At order Λ^2, parameters without any indices,

$$\Lambda^2, \quad \sigma\Lambda\Lambda. \tag{15}$$

  with indices α β,

$$\Lambda^2\ \epsilon^{\alpha\beta}, \quad \sigma\Lambda\Lambda\ \epsilon^{\alpha\beta}; \qquad \Lambda^2\ \epsilon^{\dot{\alpha}\dot{\beta}}, \quad \sigma\Lambda\Lambda\ \epsilon^{\dot{\alpha}\dot{\beta}}. \tag{16}$$

  with indices k l,

$$\Lambda^2\ \eta^{kl}, \quad \sigma\Lambda\Lambda\ \eta^{kl}, \quad \Lambda^{kl}, \quad \left(\eta\sigma\Lambda\Lambda^{k}\right)^{l}. \tag{17}$$

  with indices k $\dot{\alpha}$ β,

$$\begin{aligned} &\Lambda^2\ (\bar{\sigma}^{k})^{\dot{\alpha}\beta}, \quad \sigma\Lambda\Lambda\ (\bar{\sigma}^{k})^{\dot{\alpha}\beta}, \quad (\bar{\sigma}^{n})^{\dot{\alpha}\beta}\ \Lambda^{k}{}_{n}, \\ &\eta_{nl}\ (\bar{\sigma}^{l})^{\dot{\alpha}\beta}\ (\eta\sigma\Lambda\Lambda^{n})^{k}, \quad \eta_{nl}\ (\bar{\sigma}^{l})^{\dot{\alpha}\beta}\ (\eta\sigma\Lambda\Lambda^{k})^{n}; \end{aligned} \tag{18}$$

with indices $k\ l\ n\ o$,

$$\begin{aligned}
&\Lambda^2\ \eta^{kl}\ \eta^{no},\quad \sigma\Lambda\Lambda\ \eta^{kl}\ \eta^{no},\quad \Lambda^2\ \epsilon^{klno},\\
&\eta^{kl}\ \Lambda^{no},\quad \eta^{kl}\ (\eta\sigma\Lambda\Lambda^n)^{\,o},\quad \epsilon^{klnp}\ \Lambda_p{}^o,\quad \left(\sigma\Lambda\Lambda^{kl}\right)^{no}.
\end{aligned} \tag{19}$$

with indices $k\ l\ \alpha\ \beta$,

$$\begin{aligned}
&\Lambda^2\ \eta^{kl}\ \epsilon^{\alpha\beta},\quad \sigma\Lambda\Lambda\ \eta^{kl}\ \epsilon^{\alpha\beta},\quad \Lambda^2\ \left(\sigma^{kl}\right)^{\alpha\beta},\quad \sigma\Lambda\Lambda\ \left(\sigma^{kl}\right)^{\alpha\beta},\\
&\left(\eta\sigma\Lambda\Lambda^k\right)^{\,l}\ \epsilon^{\alpha\beta},\quad \Lambda^{kl}\ \epsilon^{\alpha\beta},\quad \Lambda^{k\alpha}\ \Lambda^{l\beta},\quad \epsilon^{klno}\ \Lambda_n{}^\alpha\ \Lambda_o{}^\beta,\\
&\left(\sigma^{kn}\right)^{\alpha\beta}\ \Lambda_n{}^l,\quad \left(\sigma\Lambda^{kn}\right)^{l\alpha}\ \Lambda_n{}^\beta,\quad \left(\eta\sigma\Lambda^k\right)^{\,\alpha}\ \Lambda^{l\beta},\quad \eta^{kl}\ (\eta\sigma\Lambda^n)^{\,\alpha}\ \Lambda_n{}^\beta.
\end{aligned} \tag{20}$$

- At order Λ^4, parameters without any indices,

$$\Lambda^2\ \Lambda^2,\quad \Lambda^2\ \sigma\Lambda\Lambda,\quad \sigma\Lambda\Lambda\ \sigma\Lambda\Lambda \tag{21}$$

with indices $\alpha\ \beta$,

$$\begin{aligned}
&\Lambda^2\ \Lambda^2\ \epsilon^{\alpha\beta},\quad \Lambda^2\ \sigma\Lambda\Lambda\ \epsilon^{\alpha\beta},\quad \sigma\Lambda\Lambda\ \sigma\Lambda\Lambda\ \epsilon^{\alpha\beta};\\
&\Lambda^2\ \Lambda^2\ \epsilon^{\dot\alpha\dot\beta},\quad \Lambda^2\ \sigma\Lambda\Lambda\ \epsilon^{\dot\alpha\dot\beta},\quad \sigma\Lambda\Lambda\ \sigma\Lambda\Lambda\ \epsilon^{\dot\alpha\dot\beta}.
\end{aligned} \tag{22}$$

with indices $k\ l$,

$$\begin{aligned}
&\Lambda^2\ \Lambda^2\ \eta^{kl},\quad \Lambda^2\ \sigma\Lambda\Lambda\ \eta^{kl},\quad \sigma\Lambda\Lambda\ \sigma\Lambda\Lambda\ \eta^{kl},\quad \Lambda^2\ \Lambda^{kl},\quad \sigma\Lambda\Lambda\ \Lambda^{kl},\\
&\Lambda^2\ \left(\eta\sigma\Lambda\Lambda^k\right)^{\,l},\quad \sigma\Lambda\Lambda\ \left(\eta\sigma\Lambda\Lambda^k\right)^{\,l},\quad (\eta\sigma\Lambda\Lambda^n)^{\,k}\ \Lambda_n{}^l.
\end{aligned} \tag{23}$$

with indices $k\ \dot\alpha\ \beta$,

$$\begin{aligned}
&\Lambda^2\ \Lambda^2\ \left(\bar\sigma^k\right)^{\dot\alpha\beta},\quad \Lambda^2\ \sigma\Lambda\Lambda\ \left(\bar\sigma^k\right)^{\dot\alpha\beta},\quad \sigma\Lambda\Lambda\ \sigma\Lambda\Lambda\ \left(\bar\sigma^k\right)^{\dot\alpha\beta},\\
&\Lambda^2\ (\bar\sigma^n)^{\dot\alpha\beta}\ \Lambda^k{}_n,\quad \sigma\Lambda\Lambda\ (\bar\sigma^n)^{\dot\alpha\beta}\ \Lambda^k{}_n,\\
&\Lambda^2\ \eta_{n_1n_2}\ (\bar\sigma^{n_2})^{\dot\alpha\beta}\ \left(\eta\sigma\Lambda\Lambda^k\right)^{n_1},\quad \Lambda^2\ \eta_{n_1n_2}\ (\bar\sigma^{n_2})^{\dot\alpha\beta}\ (\eta\sigma\Lambda\Lambda^{n_1})^k,\\
&\sigma\Lambda\Lambda\ \eta_{n_1n_2}\ (\bar\sigma^{n_2})^{\dot\alpha\beta}\ \left(\eta\sigma\Lambda\Lambda^k\right)^{n_1},\quad \sigma\Lambda\Lambda\ \eta_{n_1n_2}\ (\bar\sigma^{n_2})^{\dot\alpha\beta}\ (\eta\sigma\Lambda\Lambda^{n_1})^k,\\
&\eta_{n_1n_2}\ (\bar\sigma^{n_2})^{\dot\alpha\beta}\ (\eta\sigma\Lambda\Lambda^{n_3})^{n_1}\ \Lambda^k{}_{n_3},\quad \Lambda_{n_1n_2}\ (\bar\sigma^{n_2})^{\dot\alpha\beta}\ (\eta\sigma\Lambda\Lambda^{n_1})^k.
\end{aligned} \tag{24}$$

with indices $k\ l\ n\ o$,

$$\begin{aligned}
&\Lambda^2\ \Lambda^2\ \eta^{kl}\ \eta^{no},\quad \Lambda^2\ \sigma\Lambda\Lambda\ \eta^{kl}\ \eta^{no},\quad \sigma\Lambda\Lambda\ \sigma\Lambda\Lambda\ \eta^{kl}\ \eta^{no},\quad \Lambda^2\ \Lambda^2\ \epsilon^{klno},\\
&\Lambda^2\ \eta^{kl}\ \Lambda^{no},\quad \sigma\Lambda\Lambda\ \eta^{kl}\ \Lambda^{no},\quad \Lambda^2\ \eta^{kl}\ (\eta\sigma\Lambda\Lambda^n)^{\,o},\quad \sigma\Lambda\Lambda\ \eta^{kl}\ (\eta\sigma\Lambda\Lambda^n)^{\,o},\\
&\Lambda^2\ \epsilon^{klnp}\ \Lambda_p{}^o,\quad \Lambda^2\ \left(\sigma\Lambda\Lambda^{kl}\right)^{no},\quad \sigma\Lambda\Lambda\ \left(\sigma\Lambda\Lambda^{kl}\right)^{no},\\
&\Lambda^{kl}\ \Lambda^{no},\quad \left(\eta\sigma\Lambda\Lambda^k\right)^{\,l}\ \Lambda^{no},\quad \left(\sigma\Lambda\Lambda^{kp}\right)^{ln}\ \Lambda_p{}^o,\quad \eta^{kl}\ (\eta\sigma\Lambda\Lambda^p)^{\,n}\ \Lambda_p{}^o,\\
&\epsilon^{klpq}\ \Lambda_p{}^n\ \Lambda_q{}^o.
\end{aligned} \tag{25}$$

with indices $k\ l\ \alpha\ \beta$,

$$
\begin{aligned}
&\Lambda^2\ \Lambda^2\ \eta^{kl}\ \epsilon^{\alpha\beta},\quad \Lambda^2\ \sigma\Lambda\Lambda\ \eta^{kl}\ \epsilon^{\alpha\beta},\quad \sigma\Lambda\Lambda\ \sigma\Lambda\Lambda\ \eta^{kl}\ \epsilon^{\alpha\beta}\\
&\Lambda^2\ \Lambda^2\ \left(\sigma^{kl}\right)^{\alpha\beta},\quad \Lambda^2\ \sigma\Lambda\Lambda\ \left(\sigma^{kl}\right)^{\alpha\beta},\quad \sigma\Lambda\Lambda\ \sigma\Lambda\Lambda\left(\sigma^{kl}\right)^{\alpha\beta},\\
&\Lambda^2\ \left(\eta\sigma\Lambda\Lambda^k\right)^{l}\ \epsilon^{\alpha\beta},\quad \sigma\Lambda\Lambda\ \left(\eta\sigma\Lambda\Lambda^k\right)^{l}\ \epsilon^{\alpha\beta},\quad \Lambda^2\ \Lambda^{kl}\ \epsilon^{\alpha\beta},\quad \sigma\Lambda\Lambda\ \Lambda^{kl}\ \epsilon^{\alpha\beta},\\
&\Lambda^2\ \Lambda^{k\alpha}\ \Lambda^{l\beta},\quad \sigma\Lambda\Lambda\ \Lambda^{k\alpha}\ \Lambda^{l\beta},\quad \Lambda^2\ \epsilon^{klno}\ \Lambda_n{}^{\beta}\ \Lambda_o{}^{\alpha},\\
&\Lambda^2\ \left(\sigma^{kn}\right)^{\alpha\beta}\ \Lambda^l{}_n,\quad \sigma\Lambda\Lambda\ \left(\sigma^{kn}\right)^{\alpha\beta}\ \Lambda^l{}_n,\quad \Lambda^2\ \left(\sigma\Lambda^{kn}\right)^{l\alpha}\ \Lambda^{\beta}_n,\quad \sigma\Lambda\Lambda\ \left(\sigma\Lambda^{kn}\right)^{l\alpha}\ \Lambda^{\beta}_n,\\
&\Lambda^2\ \left(\eta\sigma\Lambda^k\right)^{\alpha}\ \Lambda^{l\beta},\quad \sigma\Lambda\Lambda\ \left(\eta\sigma\Lambda^k\right)^{\alpha}\ \Lambda^{l\beta},\quad \Lambda^2\ \eta^{kl}\ \left(\eta\sigma\Lambda^n\right)^{\alpha}\ \Lambda^{\beta}_n,\quad \sigma\Lambda\Lambda\ \eta^{kl}\ \left(\eta\sigma\Lambda^n\right)^{\alpha}\ \Lambda^{\beta}_n,\\
&\left(\eta\sigma\Lambda\Lambda^n\right)^{k}\ \Lambda^l{}_n\ \epsilon^{\alpha\beta},\quad \Lambda^k{}_n\ \Lambda^l{}_o\ \left(\sigma^{no}\right)^{\alpha\beta},\quad \Lambda^{kl}\ \left(\eta\sigma\Lambda^n\right)^{\alpha}\ \Lambda^{\beta}_n,\quad \epsilon^{knop}\ \Lambda^l{}_n\ \Lambda^{\alpha}_o\ \Lambda^{\beta}_p,\\
&\left(\sigma\Lambda\Lambda^{no}\right)^{kl}\ \Lambda^{\alpha}_n\ \Lambda^{\beta}_o,\quad \left(\eta\sigma\Lambda\Lambda^n\right)^{k}\ \Lambda^{l\alpha}\ \Lambda^{\beta}_n,\quad \left(\sigma\Lambda^{no}\right)^{k\alpha}\ \Lambda^l{}_n\ \Lambda^{\beta}_o,\quad \left(\eta\sigma\Lambda^n\right)^{\alpha}\ \Lambda^{k\beta}\ \Lambda^l{}_n. \qquad (26)
\end{aligned}
$$

4 Construct 1/2 supersymmetric invariant action

We can construct 1/2 supersymmetric invariant action by using the BFNC parameters and divergent operators as follows,

1. We can remove θ^2 in $\int d^4x\ d^4\theta\ \theta^4$ by integrating by part of D^2 .

2. We do not apply D_α, $\overline{D}_{\dot\alpha}$ on the same superfield, because we will obtain $\left(\sigma^k\right)_{\alpha\dot\beta}\ \partial_k$ by using D algebraic relations.

3. Take into account the possible simplification when combining BFNC parameters with $\partial_k\partial_l$ or $D_\alpha D_\beta$.

4. For operators with the same number of superfields and containing $D^2\overline{D}^2$, $\partial D\overline{D}$ or $\partial\partial$, we only keep the $\overline{D}^2$.

5. We define new parameters by using the BFNC parameters. At the order of Λ^2, we define X_i, $X_i{}^{k_1k_2}$, $X_i{}^{k_1k_2k_3k_4}$, $\overline{X}_i{}^{k_1k_2,k_3k_4}$, $X_i{}^{k_1\dot\alpha\beta}$, $X_i{}^{k_1k_2\alpha\beta}$, $\overline{X}_i{}^{k_1k_2\alpha\beta}$, see Eq. (A13)-(A19). At the order of Λ^4, we define Y_i, $Y_i{}^{k_1k_2}$, $\overline{Y}_i{}^{k_1k_2}$, $Y_i{}^{k_1k_2k_3k_4}$, $\overline{Y}_i{}^{k_1,k_2k_3k_4}$, $\widetilde{Y}_i{}^{k_1k_2,k_3k_4}$, $\hat{Y}_i{}^{k_1k_2,k_3k_4}$, $\widetilde{\overline{Y}}_i{}^{k_1,k_2k_3,k_4}$, $Y_i{}^{k_1\dot\alpha\beta}$, $Y_i{}^{k_1k_2\alpha\beta}$, $\overline{Y}_i{}^{k_1k_2\alpha\beta}$, $\widetilde{Y}_i{}^{k_1k_2\alpha\beta}$, see Eq. (A21)-(A32).

At the order of Λ^2, by using Eq. (1) we can construct the following action,

$$
\begin{aligned}
&\int d^4x\ d^4\theta\ \Big\{X_1\ \theta^4\ \Phi^+ + X_2\ \theta^4\left(\Phi^+\right)^2 + X_3\ \theta^4\left(\Phi^+\right)^3 + X_4\ \theta^4\left(\Phi^+\right)^4 + X_5\ \theta^4\left(\Phi^+\right)^5 + X_6\ \overline{\theta}^2\ \Phi\\
&+X_7\ \overline{\theta}^2\ \Phi\ \Phi^+ + X_8\ \overline{\theta}^2\left(D^2\Phi\right)\Phi + X_9\ \overline{\theta}^2\ \Phi\left(\Phi^+\right)^2 + X_{10}\ \overline{\theta}^2\left(D^2\Phi\right)\Phi\ \Phi^+ + X_{11}\ \overline{\theta}^2\ \Phi\left(\Phi^+\right)^3\Big\}. \qquad (27)
\end{aligned}
$$

by using Eq. (2) we obtain,

$$\int d^4x\ d^4\theta\ \Big\{ X_{12}{}^{k_1k_2}\ \theta^4\left(\partial_{k_1}\partial_{k_2}\Phi^+\right)\Phi^+ + X_{13}{}^{k_1k_2}\ \overline{\theta}^2\ \Phi\left(\partial_{k_1}\partial_{k_2}\Phi^+\right) + X_{14}{}^{k_1k_2}\ \overline{\theta}^2\left(\partial_{k_1}\partial_{k_2}D^2\Phi\right)\Phi$$
$$+X_{15}{}^{k_1k_2}\ \theta^4\left(\partial_{k_1}\partial_{k_2}\Phi^+\right)\left(\Phi^+\right)^2 + X_{16}{}^{k_1k_2}\ \overline{\theta}^2\ \Phi\left(\partial_{k_1}\Phi^+\right)\left(\partial_{k_2}\Phi^+\right) + X_{17}{}^{k_1k_2}\ \overline{\theta}^2\ \Phi\left(\partial_{k_1}\partial_{k_2}\Phi^+\right)\Phi^+$$
$$+X_{18}{}^{k_1k_2}\ \theta^4\left(\partial_{k_1}\partial_{k_2}\Phi^+\right)\left(\Phi^+\right)^3\Big\}. \tag{28}$$

by using Eq. (3) we obtain,

$$\int d^4x\ d^4\theta\ \Big\{ X_{19}{}^{k_1k_2k_3k_4}\ \theta^4\left(\partial_{k_1}\partial_{k_2}\partial_{k_3}\partial_{k_4}\Phi^+\right)\Phi^+ + X_{20}{}^{k_1k_2k_3k_4}\ \overline{\theta}^2\ \Phi\left(\partial_{k_1}\partial_{k_2}\partial_{k_3}\partial_{k_4}\Phi^+\right)$$
$$+X_{21}{}^{k_1k_2k_3k_4}\ \theta^4\left(\partial_{k_1}\partial_{k_2}\partial_{k_3}\partial_{k_4}\Phi^+\right)\left(\Phi^+\right)^2 + \overline{X}_{22}{}^{k_1k_2,k_3k_4}\ \theta^4\left(\partial_{k_1}\partial_{k_2}\Phi^+\right)\left(\partial_{k_3}\partial_{k_4}\Phi^+\right)\Phi^+\Big\}. \tag{29}$$

by using Eq. (4) we obtain,

$$\int d^4x\ d^4\theta\ \Big\{ X_{23}\ \theta^4\left(\bar{D}^2\Phi^+\right) + X_{24}\ \theta^4\left(\bar{D}^2\Phi^+\right)\Phi^+ + X_{24}\ \epsilon^{\dot\alpha\dot\beta}\ \theta^4\left(\bar{D}_{\dot\alpha}\Phi^+\right)\left(\bar{D}_{\dot\beta}\Phi^+\right)$$
$$+\frac{1}{2}\ X_{25}\ \theta^4\left(\bar{D}^2\Phi^+\right)\left(\Phi^+\right)^2 + X_{25}\ \epsilon^{\dot\alpha\dot\beta}\ \theta^4\left(\bar{D}_{\dot\alpha}\Phi^+\right)\left(\bar{D}_{\dot\beta}\Phi^+\right)\Phi^+ + \frac{1}{3}\ X_{26}\ \theta^4\left(\bar{D}^2\Phi^+\right)\left(\Phi^+\right)^3$$
$$+X_{26}\ \epsilon^{\dot\alpha\dot\beta}\ \theta^4\left(\bar{D}_{\dot\alpha}\Phi^+\right)\left(\bar{D}_{\dot\beta}\Phi^+\right)\left(\Phi^+\right)^2 + \frac{1}{4}\ X_{27}\ \theta^4\left(\bar{D}^2\Phi^+\right)\left(\Phi^+\right)^4 + X_{27}\ \epsilon^{\dot\alpha\dot\beta}\ \theta^4\left(\bar{D}_{\dot\alpha}\Phi^+\right)\left(\bar{D}_{\dot\beta}\Phi^+\right)\left(\Phi^+\right)^3$$
$$+\frac{1}{5}\ X_{28}\ \theta^4\left(\bar{D}^2\Phi^+\right)\left(\Phi^+\right)^5 + X_{28}\ \epsilon^{\dot\alpha\dot\beta}\ \theta^4\left(\bar{D}_{\dot\alpha}\Phi^+\right)\left(\bar{D}_{\dot\beta}\Phi^+\right)\left(\Phi^+\right)^4\Big\}. \tag{30}$$

by using Eq. (5) we obtain,

$$\int d^4x\ d^4\theta\ \Big\{ X_{29}\ \overline{\theta}^2\ \Phi^2 + X_{30}\ \overline{\theta}^2\ \Phi^3 + X_{31}\ \overline{\theta}^2\ \left(D^2\Phi\right)\Phi^2 + X_{32}\ \overline{\theta}^2\ (\Phi)^2\ \Phi^+ + X_{33}\ \overline{\theta}^2\ \left(D^2\Phi\right)\Phi^3$$
$$+X_{34}\ \overline{\theta}^2\ \Phi^3\ \Phi^+ + X_{35}\ \overline{\theta}^2\ \left(D^2\Phi\right)\Phi^2\ \Phi^+ + X_{36}\ \overline{\theta}^2\ \Phi^2\left(\Phi^+\right)^2 + X_{37}\ \overline{\theta}^2\ \Phi^2\left(\Phi^+\right)^3$$
$$+X_{38}\ \overline{\theta}^2\ \Phi^3\left(\Phi^+\right)^2 + X_{39}\ \overline{\theta}^2\ \left(D^2\Phi\right)\Phi^3\ \Phi^+ + X_{40}\ \overline{\theta}^2\ \Phi^3\left(\Phi^+\right)^3\Big\}. \tag{31}$$

by using Eq. (6) we obtain,

$$\int d^4x\ d^4\theta\ \Big\{ X_{41}{}^{k_1k_2}\ \overline{\theta}^2\left(\partial_{k_1}\partial_{k_2}\Phi\right)\Phi + X_{42}{}^{k_1k_2}\ \overline{\theta}^2\left(\partial_{k_1}\partial_{k_2}D^2\Phi\right)\Phi^2 + X_{43}{}^{k_1k_2}\ \overline{\theta}^2\left(D^2\Phi\right)\left(\partial_{k_1}\partial_{k_2}\Phi\right)\Phi$$
$$+X_{44}{}^{k_1k_2\alpha\beta}\ \overline{\theta}^2\left(\partial_{k_1}\partial_{k_2}D_\alpha\Phi\right)\left(D_\beta\Phi\right)\Phi + \overline{X}_{45}{}^{k_1k_2\alpha\beta}\ \overline{\theta}^2\left(\partial_{k_1}D_\alpha\Phi\right)\left(\partial_{k_2}D_\beta\Phi\right)\Phi + X_{46}\ \overline{\theta}^2\ \Phi\left(\overline{D}^2\Phi^+\right)$$
$$+X_{47}\ \overline{\theta}^2\ \Phi\left(\overline{D}^2\Phi^+\right)\Phi^+ + X_{48}\ \epsilon^{\dot\alpha\dot\beta}\ \overline{\theta}^2\ \Phi\left(\overline{D}_{\dot\alpha}\Phi^+\right)\left(\overline{D}_{\dot\beta}\Phi^+\right) + X_{49}\ \overline{\theta}^2\left(D^2\Phi\right)\Phi\left(\overline{D}^2\Phi^+\right)$$
$$+X_{50}\ \epsilon^{\alpha\beta}\ \overline{\theta}^2\left(D_\alpha\Phi\right)\left(D_\beta\Phi\right)\left(\overline{D}^2\Phi^+\right) + X_{51}{}^{k_1\dot\alpha\beta}\ \overline{\theta}^2\left(\partial_{k_1}D_\beta\Phi\right)\Phi\left(\overline{D}_{\dot\alpha}\Phi^+\right)$$
$$+X_{52}{}^{k_1\dot\alpha\beta}\ \overline{\theta}^2\left(D_\beta\Phi\right)\left(\partial_{k_1}\Phi\right)\left(\overline{D}_{\dot\alpha}\Phi^+\right) + X_{53}{}^{k_1k_2}\ \overline{\theta}^2\left(\partial_{k_1}\partial_{k_2}\Phi\right)\Phi\ \Phi^+ + X_{54}{}^{k_1k_2}\ \overline{\theta}^2\left(\partial_{k_1}\Phi\right)\left(\partial_{k_2}\Phi\right)\Phi^+$$
$$+X_{55}\ \overline{\theta}^2\ \Phi\left(\overline{D}^2\Phi^+\right)\left(\Phi^+\right)^2 + X_{56}\ \epsilon^{\dot\alpha\dot\beta}\ \overline{\theta}^2\ \Phi\left(\overline{D}_{\dot\alpha}\Phi^+\right)\left(\overline{D}_{\dot\beta}\Phi^+\right)\Phi^+ + X_{57}\ \overline{\theta}^2\left(D^2\Phi\right)\Phi\left(\overline{D}^2\Phi^+\right)\Phi^+$$
$$+X_{58}\ \epsilon^{\dot\alpha\dot\beta}\ \overline{\theta}^2\left(D^2\Phi\right)\Phi\left(\overline{D}_{\dot\alpha}\Phi^+\right)\left(\overline{D}_{\dot\beta}\Phi^+\right) + X_{59}\ \epsilon^{\alpha\beta}\ \overline{\theta}^2\left(D_\alpha\Phi\right)\left(D_\beta\Phi\right)\left(\overline{D}^2\Phi^+\right)\Phi^+$$
$$+X_{60}{}^{k_1\dot\alpha\beta}\ \overline{\theta}^2\left(\partial_{k_1}D_\beta\Phi\right)\Phi\left(\overline{D}_{\dot\alpha}\Phi^+\right)\Phi^+ + X_{61}{}^{k_1\dot\alpha\beta}\ \overline{\theta}^2\left(D_\beta\Phi\right)\left(\partial_{k_1}\Phi\right)\left(\overline{D}_{\dot\alpha}\Phi^+\right)\Phi^+$$

$$+X_{62}{}^{k_1\dot{\alpha}\beta}\ \overline{\theta}^2\left(D_\beta\Phi\right)\Phi\left(\partial_{k_1}\overline{D}_{\dot{\alpha}}\Phi^+\right)\Phi^+ + X_{63}{}^{k_1k_2}\ \overline{\theta}^2\left(\partial_{k_1}\partial_{k_2}\Phi\right)\Phi\ \left(\Phi^+\right)^2$$
$$+X_{64}{}^{k_1k_2}\ \overline{\theta}^2\left(\partial_{k_1}\Phi\right)\left(\partial_{k_2}\Phi\right)\left(\Phi^+\right)^2 + X_{65}{}^{k_1k_2}\ \overline{\theta}^2\ \Phi^2\left(\partial_{k_1}\partial_{k_2}\Phi^+\right)\Phi^+$$
$$+X_{66}\ \overline{\theta}^2\ \Phi\left(\overline{D}^2\Phi^+\right)\left(\Phi^+\right)^3 + X_{67}\ \epsilon^{\dot{\alpha}\dot{\beta}}\ \overline{\theta}^2\ \Phi\left(\overline{D}_{\dot{\alpha}}\Phi^+\right)\left(\overline{D}_{\dot{\beta}}\Phi^+\right)\left(\Phi^+\right)^2\Bigg\}. \tag{32}$$

At the order of Λ^4, we can construct the action corresponding to the operators in Eq. (7)-(11), see Eq. (A34)-(A38) in the Appendix.

5 Conclusion and outlook

In summary, at the order of Λ^2, we construct actions in Eq. (27)-(32) for divergent operators in Eq. (1)-(6). Except for the action in Eq.(30), all of the parameters are independent. After removing D^2, we found $\int d^4xd^4\theta\ \theta^4 \to \int d^4xd^4\theta\ \overline{\theta}^2$ for some terms in Eq. (27)-(30), and all terms in Eq. (31)-(32).

Because the deformed action Eq. (A4) have 1/2 supersymmetry, its effective action should also have 1/2 supersymmetry. The action in Eq. (27)-(32) that writen as a superspace integral $\int d^4xd^4\theta\ \overline{\theta}^2$ have manifest 1/2 supersymmetry, because there is no θ^2 that do not commute with the supersymmetry generator Q_α. The 1/2 supersymmetry invariance of the action with superspace integral $\int d^4xd^4\theta\ \theta^4$ in Eq. (27)-(30) can also be verified easily. At the order of Λ^2, we can add the action in Eq. (27)-(32) to the deformed action in Eq. (A4) and define a new action, which is renormalizable to all order in perturbative theory, because it contains all of the divergent operators and have 1/2 supersymmetry.

If we add the Λ^2 order action in Eq. (27)-(32) and the Λ^4 oder action in Eq. (A34)-(A38) to the deformed action in Eq. (A4), we obtain a renormalizable BFNC Wess-Zumino action at the order of Λ^4. It is straightforward to extend the construction to the order of Λ^8, which is the highest order of noncommutative parameters, because there are 8 different Fermionic noncommutative parameters $\Lambda^{k\alpha}$.

We found there are many free parameters for BFNC Wess-Zumino model and expect the introduction of supergauge symmetry [16] will reduce the number of free parameters.

Acknowledgments

The author would like to thank Professors Jian-Xin Lu, Yan-Gang Miao, Zhi-Guang Xiao for helpful discussions. This work was supported by a key grant from the NSF of China with Grant No: 11235010.

A The deformed Wess-Zumino model on BFNC superspace

The Wess-Zumino model is defined on superspace as follows,

$$\begin{aligned}\mathcal{S}_{\rm WZ} &= \int d^8z \left\{\Phi^+\Phi - \frac{m}{8}\,\Phi\left(\frac{D^2}{\Box}\Phi\right) - \frac{m^*}{8}\,\Phi^+\left(\frac{\bar{D}^2}{\Box}\Phi^+\right)\right.\\ &\qquad \left. -\frac{g}{12}\,\Phi\,\Phi\left(\frac{D^2}{\Box}\Phi\right) - \frac{g^*}{12}\,\Phi^+\,\Phi^+\left(\frac{\bar{D}^2}{\Box}\Phi^+\right)\right\}. \end{aligned} \tag{A1}$$

BFNC star product is defined as follows,

$$\mathbf{F}\star\mathbf{G} \equiv \mu\left\{\exp\left[\frac{i}{2}\,\Lambda^{k\alpha}\left(\frac{\partial}{\partial y^k}\otimes\frac{\partial}{\partial\theta^\alpha} - \frac{\partial}{\partial\theta^\alpha}\otimes\frac{\partial}{\partial y^k}\right)\right]\triangleright(\mathbf{F}\otimes\mathbf{G})\right\}, \tag{A2}$$

Taylor expansion,

$$\begin{aligned}\mathbf{F}\star\mathbf{G} &= \mathbf{FG} - \frac{i}{2}\,\Lambda^{k\alpha}\,(\partial_\alpha\mathbf{F})\,(\partial_k\mathbf{G}) + (-1)^{|\mathbf{F}|}\,\frac{i}{2}\,\Lambda^{k\alpha}\,(\partial_k\mathbf{F})\,(\partial_\alpha\mathbf{G})\\ &\quad +\frac{1}{8}\,\Lambda^{k\alpha}\,\Lambda^{l\beta}\,(\partial_k\partial_l\mathbf{F})\,(\partial_\alpha\partial_\beta\mathbf{G}) + \frac{1}{8}\,\Lambda^{k\alpha}\,\Lambda^{l\beta}\,(\partial_\alpha\partial_\beta\mathbf{F})\,(\partial_k\partial_l\mathbf{G})\\ &\quad +(-1)^{|\mathbf{F}|}\,\frac{1}{4}\,\Lambda^{k\alpha}\,\Lambda^{l\beta}\,(\partial_\beta\partial_k\mathbf{F})\,(\partial_\alpha\partial_l\mathbf{G})\\ &\quad -\frac{i}{16}\,\Lambda^{k\alpha}\,\Lambda^{l\beta}\,\Lambda^{m\zeta}\,(\partial_\alpha\partial_l\partial_m\mathbf{F})\,(\partial_\beta\partial_\zeta\partial_k\mathbf{G})\\ &\quad +(-1)^{|\mathbf{F}|}\,\frac{i}{16}\,\Lambda^{k\alpha}\,\Lambda^{l\beta}\,\Lambda^{m\zeta}\,(\partial_\alpha\partial_\beta\partial_m\mathbf{F})\,(\partial_\zeta\partial_k\partial_l\mathbf{G})\\ &\quad -\frac{1}{64}\,\Lambda^{k\alpha}\,\Lambda^{l\beta}\,\Lambda^{m\zeta}\,\Lambda^{n\iota}\,(\partial_\alpha\partial_\zeta\partial_l\partial_n\mathbf{F})\,(\partial_\beta\partial_\iota\partial_k\partial_m\mathbf{G}), \end{aligned} \tag{A3}$$

where $\partial_k \equiv \frac{\partial}{\partial y^k}$, $\partial_\alpha \equiv \frac{\partial}{\partial\theta^\alpha}$.

By replacing the ordinary products in Eq. (A1) by star products Eq. (A3), we obtain,

$$\begin{aligned}\mathcal{S}_{\rm NC} &= \int d^8z\Big\{\Phi^+\Phi - \frac{m}{8}\,\Phi\left(\frac{D^2}{\Box}\Phi\right) - \frac{m^*}{8}\,\Phi^+\left(\frac{\bar{D}^2}{\Box}\Phi^+\right)\\ &\quad -\frac{g}{12}\,\Phi\,\Phi\left(\frac{D^2}{\Box}\Phi\right) - \frac{g^*}{12}\,\Phi^+\,\Phi^+\left(\frac{\bar{D}^2}{\Box}\Phi^+\right)\\ &\quad -\frac{g}{32}\,\Lambda^{kl}\,\theta^4\,\Phi\left(D^2\Phi\right)\left(\partial_l\partial_k D^2\Phi\right) - \frac{g}{32}\,\Lambda^{kl}\,\theta^4\,\Phi\left(\partial_k D^2\Phi\right)\left(\partial_l D^2\Phi\right)\\ &\quad -\frac{g^*}{6}\,\Lambda^{kl}\,\theta^4\,\Phi^+\left(\Box\Phi^+\right)\left(\partial_k\partial_l\Phi^+\right) - \frac{g^*}{3}\,\left(\sigma\Lambda\Lambda^{kl}\right)^{no}\,\theta^4\,\Phi^+\left(\partial_k\partial_n\Phi^+\right)\left(\partial_l\partial_o\Phi^+\right)\\ &\quad +\frac{g^*}{6}\,\eta^{kl}\,\Lambda^{no}\,\theta^4\,\Phi^+\left(\partial_k\partial_n\Phi^+\right)\left(\partial_l\partial_o\Phi^+\right) - \frac{g}{16}\,\epsilon^{\alpha\beta}\,\Lambda^{kl}\,\theta^4\,\left(\partial_k D_\alpha\Phi\right)\left(\partial_l D_\beta\Phi\right)\left(D^2\Phi\right)\\ &\quad -\frac{g}{16}\,\epsilon^{\alpha\beta}\,\epsilon^{\zeta\iota}\,\Lambda^k{}_\beta\,\Lambda^l{}_\iota\,\theta^4\left(\partial_k D_\alpha\Phi\right)\left(\partial_l D_\zeta\Phi\right)\left(D^2\Phi\right)\\ &\quad -\frac{g}{3072}\,\Lambda^{kl}\,\Lambda^{no}\,\theta^4\left(D^2\Phi\right)\left(\partial_l\partial_k D^2\Phi\right)\left(\partial_o\partial_n D^2\Phi\right)\Big\}, \end{aligned} \tag{A4}$$

	dim	$U(1)_R$	$U(1)_\Phi$		dim	$U(1)_R$	$U(1)_\Phi$
m	1	0	-2	m^*	1	0	2
g	0	-1	-3	g^*	0	1	3
$(\Lambda^k{}_\alpha)^2$	-3	2	0	V	-5	2	0
$d^4\theta$	2	0	0	θ^4	-2	0	0
Φ	1	1	1	Φ^+	1	-1	-1
D_α	$\frac{1}{2}$	-1	0	$\bar{D}_{\dot\alpha}$	$\frac{1}{2}$	1	0
D^2	1	-2	0	$\bar{D}^2$	1	2	0
∂_k	1	0	0	d^4x	-4	0	0

B Determine divergent operators

We introduce two global $U(1)$ symmetries.

We obtain the following constraints on the divergent operators,

$$\begin{aligned}
\Gamma &= \int d^4x\ \lambda\ \mathcal{O}, \quad \lambda \sim \Lambda_{UV}^d\ g^{x-R}\ g^{*x} \left(\frac{m}{\Lambda_{UV}}\right)^y \left(\frac{m^*}{\Lambda_{UV}}\right)^{y+\frac{S-3R}{2}}, \\
\mathcal{O} &= d^4\theta\ (D^2)^\gamma\ (\bar{D}^2)^\delta\ (\partial D\bar{D})^\eta\ (\partial\partial)^\zeta\ V^\rho\ \Phi^\alpha\ (\Phi^+)^\beta, \quad V \equiv (\Lambda^k{}_\alpha)^2\ \theta^4, \\
d &= 2-\alpha-\beta-\gamma-\delta-2\ \zeta-2\ \eta+5\ \rho, \quad R=-\alpha+\beta+2\ \gamma-2\ \delta-2\ \rho, \quad S=-\alpha+\beta, \\
P &= d+\frac{3}{2}R-\frac{1}{2}S-2\ y = 2-2\ y-2\ \alpha+2\ \gamma-4\ \delta-2\ \zeta-2\ \eta+2\ \rho \geq 0, \\
\gamma &\leq \alpha-\eta, \quad \delta \leq \beta-\eta, \\
\gamma &\geq 0, \quad \delta \geq 0, \quad \alpha \geq 0, \quad \eta \geq 0, \quad \rho \geq 0, \quad \zeta \geq 0, \quad \beta \geq 0, \\
\rho &= 2, \\
y &\geq 0, \quad x \geq 0, \quad x+\alpha-\beta-2\ \gamma+2\ \delta+2\ \rho \geq 0, \quad y+\alpha-\beta-3\ \gamma+3\ \delta+3\ \rho \geq 0. \qquad \text{(A5)}
\end{aligned}$$

C Convention and identities

We define the following symbols,

$$\begin{aligned}
\Lambda^{kl} &\equiv \epsilon^{\alpha\beta}\ \Lambda^k{}_\beta\ \Lambda^l{}_\alpha, & \Lambda^2 &\equiv \eta_{kl}\ \Lambda^{kl}, \\
\sigma\Lambda\Lambda &\equiv \eta_{kn}\ \eta_{lo}\ \left(\sigma^{kl}\right)^{\alpha\beta}\ \Lambda^n{}_\alpha\ \Lambda^o{}_\beta, & \left(\sigma\Lambda^{kl}\right)^{n\alpha} &\equiv \left(\sigma^{kl}\right)^{\beta\alpha}\ \Lambda^n{}_\beta, \\
\left(\eta\sigma\Lambda^k\right)^\alpha &\equiv \eta_{ln}\ \left(\sigma^{nk}\right)^{\beta\alpha}\ \Lambda^l{}_\beta, & \left(\eta\sigma\Lambda\Lambda^k\right)^l &\equiv \eta_{no}\ \left(\sigma^{ok}\right)^{\alpha\beta}\ \Lambda^n{}_\alpha\ \Lambda^l{}_\beta, \\
\left(\sigma\Lambda\Lambda^{kl}\right)^{no} &\equiv \left(\sigma^{kl}\right)^{\alpha\beta}\ \Lambda^n{}_\alpha\ \Lambda^o{}_\beta. & & \qquad \text{(A6)}
\end{aligned}$$

Combine σ,

$$\sigma^l{}_{\beta\dot\gamma}\ \left(\bar\sigma^k\right)^{\dot\alpha\beta} = -\delta_{\dot\alpha\dot\gamma}\ \eta^{kl} + 2\left(\bar\sigma^{kl}\right)^{\dot\alpha}{}_{\dot\gamma},$$

$$\sigma^k{}_{\alpha\dot\beta}\ \left(\bar\sigma^l\right)^{\dot\beta\gamma} \ = \ -\delta_{\alpha\gamma}\ \eta^{kl} + 2\left(\sigma^{kl}\right)^{\gamma}_{\alpha}. \tag{A7}$$

Reduce the number of σ,

$$\begin{aligned}
\sigma^k{}_{\gamma\dot\zeta}\ \left(\bar\sigma^{lm}\right)^{\dot\zeta}{}_{\dot\alpha} &= \frac{1}{2}\ \eta^{km}\ \sigma^l{}_{\gamma\dot\alpha} - \frac{1}{2}\ \eta^{kl}\ \sigma^m{}_{\gamma\dot\alpha} - \frac{i}{2}\ \eta_{no}\ \epsilon^{oklm}\ \sigma^n{}_{\gamma\dot\alpha},\\
(\bar\sigma^m)^{\dot\beta\gamma}\ \left(\bar\sigma^{kl}\right)^{\dot\alpha}{}_{\dot\beta} &= -\frac{1}{2}\ \eta^{lm}\ \left(\bar\sigma^k\right)^{\dot\alpha\gamma} + \frac{1}{2}\ \eta^{km}\ \left(\bar\sigma^l\right)^{\dot\alpha\gamma} + \frac{i}{2}\ \eta_{no}\ \epsilon^{oklm}\ (\bar\sigma^n)^{\dot\alpha\gamma},\\
(\bar\sigma^m)^{\dot\alpha\beta}\ \left(\sigma^{kl}\right)^{\gamma}_{\beta} &= \frac{1}{2}\ \eta^{lm}\ \left(\bar\sigma^k\right)^{\dot\alpha\gamma} - \frac{1}{2}\ \eta^{km}\ \left(\bar\sigma^l\right)^{\dot\alpha\gamma} + \frac{i}{2}\ \eta_{no}\ \epsilon^{oklm}\ (\bar\sigma^n)^{\dot\alpha\gamma},\\
\sigma^m{}_{\beta\dot\gamma}\ \left(\sigma^{kl}\right)^{\beta}_{\alpha} &= -\frac{1}{2}\ \eta^{lm}\ \sigma^k{}_{\alpha\dot\gamma} + \frac{1}{2}\ \eta^{km}\ \sigma^l{}_{\alpha\dot\gamma} - \frac{i}{2}\ \eta_{no}\ \epsilon^{oklm}\ \sigma^n{}_{\alpha\dot\gamma},\\
\left(\bar\sigma^{kl}\right)^{\dot\alpha}{}_{\dot\beta}\ (\bar\sigma^{mn})^{\dot\beta}{}_{\dot\alpha} &= \frac{i}{2}\ \epsilon^{klmn} + \frac{1}{2}\ \eta^{kn}\ \eta^{lm} - \frac{1}{2}\ \eta^{km}\ \eta^{ln},\\
\left(\sigma^{kl}\right)^{\zeta}_{\alpha}\ (\sigma^{mn})^{\alpha}_{\zeta} &= -\frac{i}{2}\ \epsilon^{klmn} + \frac{1}{2}\ \eta^{kn}\ \eta^{lm} - \frac{1}{2}\ \eta^{km}\ \eta^{ln}.
\end{aligned} \tag{A8}$$

We can reduce the number of σ from 2 to 1 for the following combinations,

$$\begin{aligned}
&\left(\sigma^{kl}\right)^{\alpha\beta}\ \sigma\Lambda\Lambda^{no}{}_{kl}, \quad \eta_{kl}\ \left(\sigma^{ln}\right)^{\alpha\beta}\ (\eta\sigma\Lambda\Lambda^o)^k, \quad \eta_{kl}\ (\sigma\Lambda^{no})^{l\alpha}\ \left(\eta\sigma\Lambda^k\right)^{\beta},\\
&\eta_{ln}\ (\eta\sigma\Lambda\Lambda^n)^q\ (\sigma\Lambda\Lambda^{or})^{ls}, \quad \eta_{kl}\ \eta_{no}\ \left(\sigma\Lambda^{lq}\right)^{o\beta}\ (\sigma\Lambda^{np})^{k\alpha},\\
&(\eta\sigma\Lambda^q)^{\beta}\ (\eta\sigma\Lambda^p)^{\alpha}, \quad (\eta\sigma\Lambda\Lambda^o)^p\ (\eta\sigma\Lambda\Lambda^s)^q, \quad \sigma\Lambda\Lambda^{kl}{}_{no}\ \sigma\Lambda\Lambda^{no}{}_{kl},\\
&\eta_{kl}\ \eta_{no}\ \left(\eta\sigma\Lambda\Lambda^l\right)^o\ (\eta\sigma\Lambda\Lambda^n)^k, \quad \eta_{kl}\ \eta_{no}\ \left(\sigma\Lambda\Lambda^{lo}\right)^{np}\ (\sigma\Lambda\Lambda^{qr})^{ks},\\
&\eta_{kl}\ \eta_{no}\ \left(\sigma\Lambda\Lambda^{lp}\right)^{oq}\ (\sigma\Lambda\Lambda^{nr})^{ks}, \quad \Lambda_{kl}\ \left(\sigma\Lambda\Lambda^{kp}\right)^{qr}\ \left(\eta\sigma\Lambda\Lambda^l\right)^s,\\
&\Lambda_{kl}\ \Lambda_{no}\ \left(\sigma\Lambda\Lambda^{kn}\right)^{l_1l_2}\ \left(\sigma\Lambda\Lambda^{lo}\right)^{l_3l_4}, \quad \Lambda_{lk_1}\ \left(\sigma^{k_1k_2}\right)^{\alpha\beta}\ \left(\eta\sigma\Lambda\Lambda^l\right)^o,\\
&\Lambda_{k_1k_3}\ \eta_{k_2l_3}\ \left(\sigma^{k_1k_2}\right)^{\alpha\beta}\ \left(\sigma\Lambda\Lambda^{k_3k_4}\right)^{l_3l_4}.
\end{aligned} \tag{A9}$$

Other identities,

$$\begin{aligned}
\left(\sigma\Lambda\Lambda^{k_3k_4}\right)^{l_3l_4}\ \Lambda^{l_5l_6} &= -\left(\sigma\Lambda\Lambda^{k_3k_4}\right)^{l_6l_4}\ \Lambda^{l_3l_5} - \left(\sigma\Lambda\Lambda^{k_3k_4}\right)^{l_5l_4}\ \Lambda^{l_3l_6},\\
\Lambda^{kl}\ \Lambda_{kl} &= -\frac{1}{2}\ \Lambda^2\ \Lambda^2,\\
\left(\eta\sigma\Lambda\Lambda^k\right)^l\ \Lambda_{kl} &= -\frac{1}{2}\ \Lambda^2\ \sigma\Lambda\Lambda,\\
\Lambda^{kn}\ \Lambda^{lo}\ \eta_{kl} &= -\frac{1}{2}\ \Lambda^2\ \Lambda^{no},\\
(\eta\sigma\Lambda\Lambda^{n_1})^{k_3}\ \Lambda^{k_4}{}_{n_1}\left(\partial_{k_3}\partial_{k_4}X\right) &= -\frac{1}{2}\ \sigma\Lambda\Lambda\ \Lambda^{k_3k_4}\left(\partial_{k_3}\partial_{k_4}X\right),\\
\Lambda^{k_1k_2}\ \left(\eta\sigma\Lambda\Lambda^{k_3}\right)^{k_4}\left(\partial_{k_1}\partial_{k_2}\partial_{k_4}X\right) &= 0,\\
\Lambda^{k_1k_3}\ \Lambda^{k_2k_4}\left(\partial_{k_1}\partial_{k_2}X\right) &= \frac{1}{2}\ \Lambda^{k_1k_2}\ \Lambda^{k_3k_4}\left(\partial_{k_1}\partial_{k_2}X\right).
\end{aligned} \tag{A10}$$

D BFNC parameters

We do not allow combination of ϵ and σ, because we can use identity to cancel ϵ. For example, if we consider $\sigma\Lambda\Lambda\ \epsilon^{k_1k_2k_3k_4}$, then we have,

$$\begin{aligned}\sigma\Lambda\Lambda\ \epsilon^{k_1k_2k_3k_4} &= -2i\left(\sigma\Lambda\Lambda^{k_1k_2}\right)^{\ k_3k_4} + 2i\left(\sigma\Lambda\Lambda^{k_1k_3}\right)^{\ k_2k_4} - 2i\left(\sigma\Lambda\Lambda^{k_2k_3}\right)^{\ k_1k_4},\\ &\quad -2i\left(\sigma\Lambda\Lambda^{k_1k_4}\right)^{\ k_2k_3} + 2i\left(\sigma\Lambda\Lambda^{k_2k_4}\right)^{\ k_1k_3} - 2i\left(\sigma\Lambda\Lambda^{k_3k_4}\right)^{\ k_1k_2}.\end{aligned}$$

We also do not consider combination that contain two ϵ, because we can transform ϵ to δ by using $\epsilon^{k_1k_2k_3k_4}\ \epsilon_{l_1l_2l_3l_4} = \delta^{k_1}{}_{[l_1}\ \delta^{k_2}{}_{l_2}\ \delta^{k_3}{}_{l_3}\ \delta^{k_4}{}_{l_4]}$.

We will not use η^{kl} when constructing parameters, but we can combine η^{kl} and parameters with indices $n\ o$ to obtain parameters with indices $k\ l\ n\ o$.

Further more, we demand the number of σ can not be reduced in the combinations, we also require the combinations not producing Λ^2 or $\sigma\Lambda\Lambda$.

D.1 Λ^2

Some of the parameters with indices $k\ l$ and $k\ l\ n\ o$ can be obtained by using Λ^2 and $\sigma\Lambda\Lambda$. The parameters with indices $k\ l$ are,

$$\Lambda^2\ \eta^{kl},\quad \sigma\Lambda\Lambda\ \eta^{kl} \tag{A11}$$

with indices $k\ l\ n\ o$ are,

$$\Lambda^2\ \eta^{kl}\ \eta^{no},\quad \sigma\Lambda\Lambda\ \eta^{kl}\ \eta^{no},\quad \Lambda^2\ \epsilon^{klno}. \tag{A12}$$

To construct new parameters with indices $k\ l$ and $k\ l\ n\ o$, we consider the following combinations,

1. To construct parameters with indices $k\ l\ n\ o$ by using $\epsilon^{k_1k_2k_3k_4}\ \Lambda^{l_1\beta_1}\ \Lambda^{l_2\beta_2}$, we can contract it with $\epsilon_{\beta_1\beta_2}$ and obtain $\epsilon^{k_1k_2k_3k_4}\ \Lambda^{l_1l_2}$, where we define $\Lambda^{l_1l_2} = \Lambda^{l_1\beta_1}\ \Lambda^{l_2\beta_2}\ \epsilon_{\beta_1\beta_2}$. There are 6 indices in $\epsilon^{k_1k_2k_3k_4}\ \Lambda^{l_1l_2}$, and have to contract with a η. We can not contract k_i, because the result is 0. If we contract l_i, the result is Λ^2, which is defined as $\Lambda^2 = \eta_{kl}\ \Lambda^{kl}$. If we obtain Λ^2, then the parameters are contained in Eq. (A11), (A12). As a result we can only contract k_i and l_i in $\epsilon^{k_1k_2k_3k_4}\ \Lambda^{l_1l_2}$. Considering the symmetries of k_i and l_i, the only possible contraction is k_1 and l_1, the result is $\epsilon^{n_1k_2k_3k_4}\ \Lambda_{n_1}{}^{l_2}$. We can only obtain parameters with indices $k\ l\ n\ o$ by using $\epsilon^{k_1k_2k_3k_4}\ \Lambda^{l_1\beta_1}\ \Lambda^{l_2\beta_2}$. If we want to construct parameters with indices $k\ l$, then we must contract $\epsilon^{n_1k_2k_3k_4}\ \Lambda_{n_1}{}^{l_2}$ with a η, due to the antisymmetric indices of ϵ, the result is 0.

2. Starting from $\Lambda^{l_1\beta_1}\ \Lambda^{l_2\beta_2}$, if we contract it with $\epsilon_{\beta_1\beta_2}$, we will obtain $\Lambda^{l_1l_2}$. If we combine $\Lambda^{l_1l_2}$ and $\eta^{l_3l_4}$, the result is $\Lambda^{l_1l_2}\ \eta^{l_3l_4}$.

3. For $\left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}\ \Lambda^{l_1\beta_1}\ \Lambda^{l_2\beta_2}$, if we contract it with $\epsilon_{\beta_1\beta_2}$, the result have symmetric indices $\alpha_1\alpha_2$. If we contract $\alpha_1\alpha_2$ with $\epsilon_{\alpha_1\alpha_2}$, the result is 0. So we can only contract $\left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}\ \Lambda^{l_1\beta_1}\ \Lambda^{l_2\beta_2}$ with $\epsilon_{\alpha_1\beta_1}\ \epsilon_{\alpha_2\beta_2}$ and obtain $\left(\sigma\Lambda\Lambda^{k_1k_2}\right)^{l_1l_2}$. We define $\left(\sigma\Lambda\Lambda^{k_1k_2}\right)^{l_1l_2} = \left(\sigma^{k_1k_2}\right)^{\alpha\beta}\ \Lambda^{l_1}{}_{\alpha}\ \Lambda^{l_2}{}_{\beta}$ (where k_1, k_2 are antisymmetric, l_1, l_2 are also antisymmetric). We also define $\Lambda^{l}{}_{\alpha} = \epsilon_{\alpha\beta}\ \Lambda^{l\beta}$. $\left(\sigma\Lambda\Lambda^{k_1k_2}\right)^{l_1l_2}$ contain indices $k\ l\ n\ o$, we can obtain parameters with indices $k\ l$ by contracting it with $\eta_{k_1l_1}$, the result is $\left(\eta\sigma\Lambda\Lambda^{k_2}\right)^{l_2}$. We define $\left(\eta\sigma\Lambda\Lambda^{k_2}\right)^{l_2} = \eta_{k_1l_1}\ \left(\sigma\Lambda\Lambda^{k_1k_2}\right)^{l_1l_2}$. We can construct parameters with indices $k\ l\ n\ o$ by using $\left(\eta\sigma\Lambda\Lambda^{k_2}\right)^{l_2}$, that is $\left(\eta\sigma\Lambda\Lambda^{k_2}\right)^{l_2}\ \eta^{l_3l_4}$.

4. Let us consider $\left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}\ \left(\sigma^{k_3k_4}\right)^{\alpha_3\alpha_4}\ \Lambda^{l_1\beta_1}\ \Lambda^{l_2\beta_2}$. We have to contract $\alpha_1\ \alpha_2\ \alpha_3\ \alpha_4\ \beta_1\ \beta_2$ by using ϵ. The indices α_i can not be contracted with ϵ, otherwise the number of σ will be reduced from 2 to 1. We have already considered the combination with one σ. If we contract β_i with ϵ, then we must also contract α_i with ϵ. Anyway, we have to contract α_i. So this combination do not give new parameters.

5. The combinations with more σ such as $\left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}\ \left(\sigma^{k_3k_4}\right)^{\alpha_3\alpha_4}\ \left(\sigma^{k_5k_6}\right)^{\alpha_5\alpha_6}\ \Lambda^{l_1\beta_1}\ \Lambda^{l_2\beta_3}$ will not give new parameters.

At the end we obtain all of the parameters with indices $k\ l$ and $k\ l\ n\ o$, see Eq. (17), (19).

In the following, we construct parameters with indices $k\ l\ \alpha\ \beta$ from parameters with indices $k\ l$ or $k\ l\ n\ o$.

1. Combine $\epsilon^{\alpha_1\alpha_2}$ and parameters with indices $k\ l$, we obtain $\left(\eta\sigma\Lambda\Lambda^{k_1}\right)^{l_1}\ \epsilon^{\alpha_1\alpha_2}$, $\Lambda^{l_1l_2}\ \epsilon^{\alpha_1\alpha_2}$, $\Lambda^2\ \eta^{k_1k_2}\ \epsilon^{\alpha_1\alpha_2}$ and $\sigma\Lambda\Lambda\ \eta^{k_1k_2}\ \epsilon^{\alpha_1\alpha_2}$.

2. Combine $\left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}$ with parameters without indices, we obtain $\Lambda^2\ \left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}$ and $\sigma\Lambda\Lambda\ \left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}$.

3. Combine $\left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}$ and parameters with indices $k\ l$, we have 3 possible forms,

   (a) $\Lambda^2\ \left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}\ \eta^{l_1l_2}$ and $\sigma\Lambda\Lambda\ \left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}\ \eta^{l_1l_2}$ do not give new parameters.

   (b) From $\left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}\ \Lambda^{l_1l_2}$ we can obtain $\left(\sigma^{n_1k_2}\right)^{\alpha_1\alpha_2}\ \Lambda_{n_1}{}^{l_2}$.

   (c) We can reduce the number of σ in $\left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}\ \left(\eta\sigma\Lambda\Lambda^{k_3}\right)^{l_3}$ by using identities.

4. Now consider combining $\left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}$ with parameters with indices $k\ l\ n\ o$:

   (a) We can cancel $\epsilon^{l_1l_2l_3l_4}$ in $\Lambda^2\ \left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}\ \epsilon^{l_1l_2l_3l_4}$.

   (b) $\Lambda^2\ \left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}\ \eta^{l_1l_2}\ \eta^{l_3l_4}$ and $\sigma\Lambda\Lambda\ \left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}\ \eta^{l_1l_2}\ \eta^{l_3l_4}$ do not give new parameters.

   (c) The $\epsilon^{n_1l_1l_2l_3}$ in $\left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}\ \epsilon^{n_1l_1l_2l_3}\ \Lambda_{n_1}{}^{l_4}$ can be removed.

   (d) For $\left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}\ \left(\sigma\Lambda\Lambda^{k_3k_4}\right)^{l_3l_4}$, if we contract k_1, k_2 with k_3, k_4, then we can reduce the number of σ. If we contract k_1, k_2 with l_3, l_4, we can also reduce the number of σ by using identities.

   (e) We can reduce the number of σ in $\left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}\ \eta^{l_1l_2}\ \left(\eta\sigma\Lambda\Lambda^{k_3}\right)^{l_3}$.

   (f) $\left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}\ \eta^{l_1l_2}\ \Lambda^{l_3l_4}$ do not give new parameters.

   We conclude that no new parameters will be produced by combing $\left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}$ with parameters with indices $k\ l\ n\ o$.

We found that we only need to construct parameters with indices $k\ l\ \alpha\ \beta$ by using at most one $\left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}$, at most one $\epsilon^{k_1k_2k_3k_4}$, one $\Lambda^{l_1\beta_1}$ and one $\Lambda^{l_2\beta_2}$. We list them as follows,

1. Without $\epsilon^{k_1k_2k_3k_4}$ and $\left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}$ we obtain $\Lambda^{l_1\beta_1}\ \Lambda^{l_2\beta_2}$.

2. Form $\epsilon^{k_1k_2k_3k_4}\ \Lambda^{l_1\beta_1}\ \Lambda^{l_2\beta_2}$ we obtain $\epsilon^{k_1k_2n_1n_2}\ \Lambda_{n_1}{}^{\beta_1}\ \Lambda_{n_2}{}^{\beta_2}$.

3. Contract $\left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}\ \Lambda^{l_1\beta_1}\ \Lambda^{l_2\beta_2}$ with $\epsilon_{\beta_1\beta_2}$, we obtain $\left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}\ \Lambda^{l_1l_2}$. Contract it with $\eta_{k_1l_1}$, the result is $\left(\sigma^{n_1k_2}\right)^{\alpha_1\alpha_2}\ \Lambda_{n_1}{}^{l_2}$.

4. Contract $\left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}\ \Lambda^{l_1\beta_1}\ \Lambda^{l_2\beta_2}$ with $\epsilon_{\alpha_1\beta_1}$, the result is $\left(\sigma\Lambda^{k_1k_2}\right)^{l_1\alpha_2}\ \Lambda^{l_2\beta_2}$, then we have 2 possibilities:

   (a) Contract $\left(\sigma\Lambda^{k_1k_2}\right)^{l_1\alpha_2}\ \Lambda^{l_2\beta_2}$ with $\eta_{k_1l_1}$, we obtain $\left(\eta\sigma\Lambda^{k_1}\right)^{\alpha_2}\ \Lambda^{l_2\beta_2}$. Contract it with $\eta_{k_1l_2}$, the result is $(\eta\sigma\Lambda^{n_1})^{\alpha_1}\ \Lambda_{n_1}{}^{\beta_2}$. By combining it with $\eta^{k_1k_2}$ we obtain $\eta^{k_1k_2}\ (\eta\sigma\Lambda^{n_1})^{\alpha_1}\ \Lambda_{n_1}{}^{\beta_2}$.

   (b) Contract $\left(\sigma\Lambda^{k_1k_2}\right)^{l_1\alpha_2}\ \Lambda^{l_2\beta_2}$ with $\eta_{k_2l_2}$, the result is $\left(\sigma\Lambda^{n_1k_2}\right)^{l_1\alpha_2}\ \Lambda_{n_1}{}^{\beta_2}$.

At the end we obtain all possible parameters with indices $k\ l\ \alpha\ \beta$, see Eq. (20).

D.2 Λ^4

In the following we will construct parameters with indices $k\ l$ and $k\ l\ n\ o$.

1. From $\Lambda^{l_1\beta_1}\ \Lambda^{l_2\beta_2}\ \Lambda^{l_3\beta_3}\ \Lambda^{l_4\beta_4}$ we obtain $\Lambda^{l_1l_2}\ \Lambda^{l_3l_4}$.

2. From $\epsilon^{k_1k_2k_3k_4}\ \Lambda^{l_1\beta_1}\ \Lambda^{l_2\beta_2}\ \Lambda^{l_3\beta_3}\ \Lambda^{l_4\beta_4}$ we have $\epsilon^{k_1k_2k_3k_4}\ \Lambda^{l_1l_2}\ \Lambda^{l_3l_4}$. By contracting it with $\eta_{k_1l_1}\ \eta_{k_2l_3}$, we obtain $\epsilon^{n_1n_2k_3k_4}\ \Lambda_{n_1}{}^{l_2}\ \Lambda_{n_2}{}^{l_4}$.

3. Contract $\left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}\ \Lambda^{l_1\beta_1}\ \Lambda^{l_2\beta_2}\ \Lambda^{l_3\beta_3}\ \Lambda^{l_4\beta_4}$ with $\epsilon_{\alpha_1\beta_1}\ \epsilon_{\alpha_2\beta_2}\ \epsilon_{\beta_3\beta_4}$, the result is $\left(\sigma\Lambda\Lambda^{k_1k_2}\right)^{l_1l_2}\ \Lambda^{l_3l_4}$. If we contract l_i, we will obtain Λ^2. So we need to consider the following two cases,

   (a) Contract $\left(\sigma\Lambda\Lambda^{k_1k_2}\right)^{l_1l_2}\ \Lambda^{l_3l_4}$ with $\eta_{k_1l_1}$ we have $\left(\eta\sigma\Lambda\Lambda^{k_1}\right)^{l_1}\ \Lambda^{l_3l_4}$. Then contract it with $\eta_{k_1l_3}$, the result is $\left(\eta\sigma\Lambda\Lambda^{n_1}\right)^{l_1}\ \Lambda_{n_1}{}^{l_4}$. The other contractions will generate Λ^2 or $\sigma\Lambda\Lambda$.

   (b) Contract $\left(\sigma\Lambda\Lambda^{k_1k_2}\right)^{l_1l_2}\ \Lambda^{l_3l_4}$ with $\eta_{k_1l_3}$ we obtain $\left(\sigma\Lambda\Lambda^{n_1k_2}\right)^{l_1l_2}\ \Lambda_{n_1}{}^{l_4}$, where k_2 can be contracted with $l_1,\ l_2,\ l_4$. If we contract k_2 with l_4 by multiplying $\eta_{k_2l_4}$, the result is 0. If we contract k_2 with l_1 or l_2, the result is $\left(\eta\sigma\Lambda\Lambda^{n_1}\right)^{l_2}\ \Lambda_{n_1}{}^{l_4}$. At the end we obtain only one parameter with indices kl, that is $\left(\eta\sigma\Lambda\Lambda^{n_1}\right)^{l_1}\ \Lambda_{n_1}{}^{l_2}$. Contract it with $\eta^{k_1k_2}$ we obtain parameter with indices $k\ l\ n\ o$, that is $\eta^{k_1k_2}\ \left(\eta\sigma\Lambda\Lambda^{n_1}\right)^{l_1}\ \Lambda_{n_1}{}^{l_2}$.

4. Consider $\left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}\ \left(\sigma^{k_3k_4}\right)^{\alpha_3\alpha_4}\ \Lambda^{l_1\beta_1}\ \Lambda^{l_2\beta_2}\ \Lambda^{l_3\beta_3}\ \Lambda^{l_4\beta_4}$, if we contract $\epsilon_{\alpha_1\beta_1}\ \epsilon_{\alpha_2\beta_2}\ \epsilon_{\alpha_3\beta_3}\ \epsilon_{\alpha_4\beta_4}$ with it, the result is $\left(\sigma\Lambda\Lambda^{k_1k_2}\right)^{l_1l_2}\ \left(\sigma\Lambda\Lambda^{k_3k_4}\right)^{l_3l_4}$. To construct parameters with indices $k\ l\ n\ o$, we should contract it with two η. k_i can not be contracted. l_i can not be contacted too. Further more, if we contract k_i in $\left(\sigma\Lambda\Lambda^{k_1k_2}\right)^{l_1l_2}$ with l_i in $\left(\sigma\Lambda\Lambda^{k_3k_4}\right)^{l_3l_4}$, we can transform it to contraction of k_i with l_j, where l_j is in $\left(\sigma\Lambda\Lambda^{k_1k_2}\right)^{l_1l_2}$, by using Fierz identities,

$$
\begin{aligned}
& \eta_{k_1l_4}\ \left(\sigma\Lambda\Lambda^{k_1k_2}\right)^{l_1l_2}\ \left(\sigma\Lambda\Lambda^{k_3k_4}\right)^{l_3l_4} \\
= \ & \eta_{k_1l_4}\ \left(\sigma\Lambda^{k_1k_2}\right)^{l_1\alpha}\ \Lambda^{l_2}{}_{\alpha}\ \left(\sigma\Lambda^{k_3k_4}\right)^{l_3\beta}\ \Lambda^{l_4}{}_{\beta} \\
= \ & -\eta_{k_1l_4}\ \left(\sigma\Lambda^{k_1k_2}\right)^{l_1\alpha}\ \left(\sigma\Lambda^{k_3k_4}\right)^{l_3\beta}\ \Lambda^{l_2}{}_{\alpha}\ \Lambda^{l_4}{}_{\beta} \\
= \ & -\eta_{k_1l_4}\ \left(\sigma\Lambda^{k_1k_2}\right)^{l_1\alpha}\ \left(\sigma\Lambda^{k_3k_4}\right)^{l_3\beta}\ \left(\Lambda^{l_2}{}_{\beta}\ \Lambda^{l_4}{}_{\alpha}+\epsilon_{\alpha\beta}\ \Lambda^{l_2l_4}\right) \\
= \ & -\eta_{k_1l_4}\ \left(\sigma\Lambda\Lambda^{k_1k_2}\right)^{l_1l_4}\ \left(\sigma\Lambda\Lambda^{k_3k_4}\right)^{l_3l_2}-\eta_{k_1l_4}\ \left(\sigma\Lambda^{k_1k_2}\right)^{l_1\alpha}\ \left(\sigma\Lambda^{k_3k_4}\right)^{l_3\beta}\ \epsilon_{\alpha\beta}\ \Lambda^{l_2l_4} \\
= \ & \left(\eta\sigma\Lambda\Lambda^{k_2}\right)^{l_1}\ \left(\sigma\Lambda\Lambda^{k_3k_4}\right)^{l_3l_2}+X.
\end{aligned}
$$

   where X represents expression in which the number of σ is reduced. So we only need to contract $\left(\sigma\Lambda\Lambda^{k_1k_2}\right)^{l_1l_2}\ \left(\sigma\Lambda\Lambda^{k_3k_4}\right)^{l_3l_4}$ with $\eta_{k_2l_2}$, the result is $\left(\eta\sigma\Lambda\Lambda^{k_1}\right)^{l_1}\ \left(\sigma\Lambda\Lambda^{k_3k_4}\right)^{l_3l_4}$. We need to contract it with a η. There are 2 possibilities,

   (a) Contract $\left(\eta\sigma\Lambda\Lambda^{k_1}\right)^{l_1}\ \left(\sigma\Lambda\Lambda^{k_3k_4}\right)^{l_3l_4}$ with $\eta_{k_1l_1}$, we obtain $\sigma\Lambda\Lambda$. We do not consider this case.

   (b) Contract $\left(\eta\sigma\Lambda\Lambda^{k_1}\right)^{l_1}\ \left(\sigma\Lambda\Lambda^{k_3k_4}\right)^{l_3l_4}$ with $\eta_{k_3l_3}$, the result is $\left(\eta\sigma\Lambda\Lambda^{k_1}\right)^{l_1}\ \left(\eta\sigma\Lambda\Lambda^{k_4}\right)^{l_4}$, we can reduce the number of σ by using identities.

In summary, $\left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}\ \left(\sigma^{k_3k_4}\right)^{\alpha_3\alpha_4}\ \Lambda^{l_1\beta_1}\ \Lambda^{l_2\beta_2}\ \Lambda^{l_3\beta_3}\ \Lambda^{l_4\beta_4}$ do not generate new parameters.

5. Consider $\left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}\ \left(\sigma^{k_3k_4}\right)^{\alpha_3\alpha_4}\ \left(\sigma^{k_5k_6}\right)^{\alpha_5\alpha_6}\ \Lambda^{l_1\beta_1}\ \Lambda^{l_2\beta_2}\ \Lambda^{l_3\beta_3}\ \Lambda^{l_4\beta_4}$. Because the number of α_i is 6, the number of β_i is 4, at least 2 α_i have to be contracted with ϵ. Then we can reduce the number of σ and do not obtain new parameters.

At the end, we obtain all parameters with indices $k\ l$ and $k\ l\ n\ o$, see Eq. (23), (25).

In the following we will construct parameters with indices $k\ l\ \alpha\ \beta$ from parameters with indices $k\ l$ or $k\ l\ n\ o$. Some of them can be obtained by combine Λ^2 or $\sigma\Lambda\Lambda$ and the parameters with indices $k\ l\ \alpha\ \beta$ at order Λ^2. The others are new parameters. We list the them as follows,

1. By combining $\epsilon^{\alpha_1\beta_1}$ and parameters with indices $k\ l$, we obtain $\epsilon^{\alpha_1\beta_1}\ \left(\eta\sigma\Lambda\Lambda^{n_1}\right)^{l_1}\ {\Lambda_{n_1}}^{l_2}$.
2. Combining $\left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}$ and parameters without indices do not give new parameters.
3. By combining $\left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}$ and parameters with indices $k\ l$, we obtain $\left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}\ \left(\eta\sigma\Lambda\Lambda^{n_1}\right)^{l_1}\ {\Lambda_{n_1}}^{l_2}$, it do not give new parameters.
4. By combining $\left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}$ and parameters with indices $k\ l\ n\ o$, we have 3 possibilities,
   (a) From $\left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}\ \Lambda^{l_1l_2}\ \Lambda^{l_3l_4}$ we have $\left(\sigma^{n_1n_2}\right)^{\alpha_1\alpha_2}\ {\Lambda_{n_1}}^{l_2}\ {\Lambda_{n_2}}^{l_4}$.
   (b) In $\left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}\ \epsilon^{n_1n_2k_3k_4}\ {\Lambda_{n_1}}^{l_1}\ {\Lambda_{n_2}}^{l_2}$ we can remove $\epsilon^{n_1n_2k_3k_4}$ and do not obtain new parameters.
   (c) For $\left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}\ \left(\eta\sigma\Lambda\Lambda^{k_3}\right)^{l_3}\ \Lambda^{l_1l_2}$, $\left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}\ \left(\sigma\Lambda\Lambda^{n_1k_3}\right)^{l_1l_2}\ {\Lambda_{n_1}}^{l_3}$ and $\left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}\ \eta^{k_3k_4}\ \left(\eta\sigma\Lambda\Lambda^{n_1}\right)^{l_1}\ {\Lambda_{n_1}}^{l_2}$, we can reduce the number of σ in them and do not obtain new parameters.

In summary, we only need to use at most one $\left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}$, one $\epsilon^{k_1k_2k_3k_4}$ and $\Lambda^{l_1\beta_1}\ \Lambda^{l_2\beta_2}\ \Lambda^{l_3\beta_3}\ \Lambda^{l_4\beta_4}$ to construct parameters with indices $k\ l\ \alpha\ \beta$, we have 2 possibilities,

1. Due to different contractions of $\left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}\ \Lambda^{l_1\beta_1}\ \Lambda^{l_2\beta_2}\ \Lambda^{l_3\beta_3}\ \Lambda^{l_4\beta_4}$ with ϵ, we can obtain 3 forms: $\left(\sigma^{k_1k_2}\right)^{\alpha_1\alpha_2}\ \Lambda^{l_1l_2}\ \Lambda^{l_3l_4}$, $\left(\sigma\Lambda^{k_1k_2}\right)^{l_1\alpha_2}\ \Lambda^{l_2\beta_2}\ \Lambda^{l_3l_4}$, $\left(\sigma\Lambda\Lambda^{k_1k_2}\right)^{l_1l_2}\ \Lambda^{l_3\beta_3}\ \Lambda^{l_4\beta_4}$. By contracting them with η, we obtain 6 parameters,

$$\left(\sigma^{n_1n_2}\right)^{\alpha_1\alpha_2}\ {\Lambda^{l_2}}_{n_1}\ {\Lambda^{l_4}}_{n_2},\quad \left(\sigma\Lambda\Lambda^{n_1n_2}\right)^{l_1l_2}\ {\Lambda_{n_1}}^{\beta_3}\ {\Lambda_{n_2}}^{\beta_4},\quad \left(\eta\sigma\Lambda\Lambda^{n_1}\right)^{l_2}\ {\Lambda_{n_1}}^{\beta_3}\ \Lambda^{l_4\beta_4}.$$

$$\left(\eta\sigma\Lambda^{n_1}\right)^{\alpha_2}\ \Lambda^{l_2\beta_2}\ {\Lambda^{l_4}}_{n_1},\quad \left(\eta\sigma\Lambda^{n_1}\right)^{\alpha_2}\ {\Lambda_{n_1}}^{\beta_2}\ \Lambda^{l_3l_4},\quad \left(\sigma\Lambda^{n_1n_2}\right)^{l_3\alpha_2}\ {\Lambda^{l_4}}_{n_2}\ {\Lambda_{n_1}}^{\beta_2},$$

2. If we use $\epsilon^{k_1k_2k_3k_4}$, the result is $\epsilon^{k_1n_1n_2n_3}\ {\Lambda^{l_1}}_{n_1}\ {\Lambda_{n_2}}^{\beta_3}\ {\Lambda_{n_3}}^{\beta_4}$.

At the end we obtain all parameters with indices $k\ l\ \alpha\ \beta$, see Eq. (26).

E The 1/2 supersymmetry invariant action

E.1 Λ^2

We define,

$$
\begin{aligned}
X_i &= \Lambda^2\ x_{i,1} + \sigma\Lambda\Lambda\ x_{i,2}, && \text{(A13)}\\
X_i{}^{k_1k_2} &= x_{i,1}\ \left(\eta\sigma\Lambda\Lambda^{k_1}\right)^{k_2} + x_{i,2}\ \eta^{k_1k_2}\ \Lambda^2 + x_{i,3}\ \Lambda^{k_1k_2} + x_{i,4}\ \eta^{k_1k_2}\ \sigma\Lambda\Lambda, && \text{(A14)}\\
X_i{}^{k_1k_2k_3k_4} &= x_{i,1}\ \eta^{k_1k_2}\ \left(\eta\sigma\Lambda\Lambda^{k_3}\right)^{k_4} + x_{i,2}\ \eta^{k_1k_2}\ \eta^{k_3k_4}\ \Lambda^2 + x_{i,3}\ \eta^{k_3k_4}\ \Lambda^{k_1k_2}\\
&\quad + x_{i,4}\ \eta^{k_1k_2}\ \eta^{k_3k_4}\ \sigma\Lambda\Lambda, && \text{(A15)}\\
\overline{X}_i{}^{k_1k_2,k_3k_4} &= x_{i,1}\ \eta^{k_1k_2}\ \left(\eta\sigma\Lambda\Lambda^{k_3}\right)^{k_4} + x_{i,2}\ \eta^{k_1k_3}\ \left(\eta\sigma\Lambda\Lambda^{k_2}\right)^{k_4} + x_{i,3}\ \eta^{k_1k_3}\ \eta^{k_2k_4}\ \Lambda^2\\
&\quad + x_{i,4}\ \eta^{k_1k_2}\ \eta^{k_3k_4}\ \Lambda^2 + x_{i,5}\ \eta^{k_1k_2}\ \Lambda^{k_3k_4} + x_{i,6}\ \eta^{k_1k_3}\ \Lambda^{k_2k_4}\\
&\quad + x_{i,7}\ \eta^{k_1k_3}\ \eta^{k_2k_4}\ \sigma\Lambda\Lambda + x_{i,8}\ \eta^{k_1k_2}\ \eta^{k_3k_4}\sigma\Lambda\Lambda + x_{i,9}\ \left(\sigma\Lambda\Lambda^{k_1k_3}\right)^{k_2k_4}, && \text{(A16)}\\
X_i{}^{k_1\dot\alpha\beta} &= x_{i,1}\ \eta_{k_2k_3}\ \left(\overline{\sigma}^{k_3}\right)^{\dot\alpha\beta}\ \left(\eta\sigma\Lambda\Lambda^{k_1}\right)^{k_2} + x_{i,2}\ \eta_{k_2k_3}\ \left(\overline{\sigma}^{k_3}\right)^{\dot\alpha\beta}\ \left(\eta\sigma\Lambda\Lambda^{k_2}\right)^{k_1}\\
&\quad + x_{i,3}\ \Lambda^2\ \left(\overline{\sigma}^{k_1}\right)^{\dot\alpha\beta} + x_{i,4}\ \sigma\Lambda\Lambda\ \left(\overline{\sigma}^{k_1}\right)^{\dot\alpha\beta} + x_{i,5}\ \left(\overline{\sigma}^{k_2}\right)^{\dot\alpha\beta}\ \Lambda^{k_1}{}_{k_2}, && \text{(A17)}\\
X_i{}^{k_1k_2\alpha\beta} &= x_{i,1}\ \eta^{k_1k_2}\ \epsilon^{\alpha\beta}\ \Lambda^2 + x_{i,2}\ \eta^{k_1k_2}\ \epsilon^{\alpha\beta}\ \sigma\Lambda\Lambda + x_{i,3}\ \epsilon^{\alpha\beta}\ \left(\eta\sigma\Lambda\Lambda^{k_1}\right)^{k_2}\\
&\quad + x_{i,4}\ \epsilon^{\alpha\beta}\ \Lambda^{k_1k_2} + x_{i,5}\ \left(\sigma^{k_1k_3}\right)^{\alpha\beta}\ \Lambda^{k_2}{}_{k_3} + x_{i,6}\ \left(\sigma\Lambda^{k_1k_3}\right)^{k_2\alpha}\ \Lambda_{k_3}{}^{\beta}\\
&\quad + x_{i,7}\ \left(\eta\sigma\Lambda^{k_1}\right)^{\alpha}\ \Lambda^{k_2\beta} + x_{i,8}\ \eta^{k_1k_2}\ \left(\eta\sigma\Lambda^{k_3}\right)^{\alpha}\ \Lambda_{k_3}{}^{\beta}, && \text{(A18)}\\
\overline{X}_i{}^{k_1k_2\alpha\beta} &= x_{i,1}\ \eta^{k_1k_2}\ \epsilon^{\alpha\beta}\ \Lambda^2 + x_{i,2}\ \eta^{k_1k_2}\ \epsilon^{\alpha\beta}\ \sigma\Lambda\Lambda + x_{i,3}\ \Lambda^2\ \left(\sigma^{k_1k_2}\right)^{\alpha\beta}\\
&\quad + x_{i,4}\ \sigma\Lambda\Lambda\ \left(\sigma^{k_1k_2}\right)^{\alpha\beta} + x_{i,5}\ \epsilon^{\alpha\beta}\ \left(\eta\sigma\Lambda\Lambda^{k_1}\right)^{k_2} + x_{i,6}\ \epsilon^{\alpha\beta}\ \Lambda^{k_1k_2}\\
&\quad + x_{i,7}\ \Lambda^{k_1\alpha}\ \Lambda^{k_2\beta} + x_{i,8}\ \epsilon^{k_1k_2k_3k_4}\ \Lambda_{k_3}{}^{\alpha}\ \Lambda_{k_4}{}^{\beta} + x_{i,9}\ \left(\sigma^{k_1k_3}\right)^{\alpha\beta}\ \Lambda^{k_2}{}_{k_3}\\
&\quad + x_{i,10}\ \left(\sigma\Lambda^{k_1k_3}\right)^{k_2\alpha}\ \Lambda_{k_3}{}^{\beta} + x_{i,11}\ \left(\eta\sigma\Lambda^{k_1}\right)^{\alpha}\ \Lambda^{k_2\beta}. && \text{(A19)}
\end{aligned}
$$

where $x_{i,j}$ are free parameters. They can combine with operators as follows,

$$
\begin{aligned}
&X_i{}^{k_1k_2}\ \Phi\ \left(\partial_{k_1}\partial_{k_2}\Phi^+\right), \quad X_i{}^{k_1k_2k_3k_4}\ \Phi\ \left(\partial_{k_1}\partial_{k_2}\partial_{k_3}\partial_{k_4}\Phi^+\right),\\
&\overline{X}_i{}^{k_1k_2,k_3k_4}\ \left(\partial_{k_1}\partial_{k_2}\Phi^+\right)\left(\partial_{k_3}\partial_{k_4}\Phi^+\right)\Phi^+, \quad X_i{}^{k_1\dot\alpha\beta}\ \left(D_\beta\Phi\right)\left(\partial_{k_1}\Phi\right)\left(\overline{D}_{\dot\alpha}\Phi^+\right)\\
&X_i{}^{k_1k_2\alpha\beta}\ \left(\partial_{k_1}\partial_{k_2}D_\alpha\Phi\right)\left(D_\beta\Phi\right)\Phi, \quad \overline{X}_i{}^{k_1k_2\alpha\beta}\left(\partial_{k_1}D_\alpha\Phi\right)\left(\partial_{k_2}D_\beta\Phi\right)\Phi. && \text{(A20)}
\end{aligned}
$$

E.2 Λ^4

We define,

$$
Y_i = y_{i,1}\ \Lambda^2\ \Lambda^2 + y_{i,2}\ \Lambda^2\ \sigma\Lambda\Lambda + y_{i,3}\ \sigma\Lambda\Lambda\ \sigma\Lambda\Lambda, \qquad \text{(A21)}
$$

$$
\begin{aligned}
Y_i^{k_1k_2} &= y_{i,1}\,\eta^{k_1k_2}\,\Lambda^2\,\Lambda^2 + y_{i,2}\,\eta^{k_1k_2}\,\Lambda^2\,\sigma\Lambda\Lambda + y_{i,3}\,\eta^{k_1k_2}\,\sigma\Lambda\Lambda\,\sigma\Lambda\Lambda \\
&\quad + y_{i,4}\,\Lambda^2\,\Lambda^{k_1k_2} + y_{i,5}\,\sigma\Lambda\Lambda\,\Lambda^{k_1k_2} + y_{i,6}\,\Lambda^2\left(\eta\sigma\Lambda\Lambda^{k_1}\right)^{k_2} \\
&\quad + y_{i,7}\,\sigma\Lambda\Lambda\,\left(\eta\sigma\Lambda\Lambda^{k_1}\right)^{k_2}, \qquad \text{(A22)} \\
\overline{Y}_i^{k_1k_2} &= y_{i,1}\,\eta^{k_1k_2}\,\Lambda^2\,\Lambda^2 + y_{i,2}\,\eta^{k_1k_2}\,\Lambda^2\,\sigma\Lambda\Lambda + y_{i,3}\,\eta^{k_1k_2}\,\sigma\Lambda\Lambda\,\sigma\Lambda\Lambda \\
&\quad + y_{i,4}\,\Lambda^2\,\Lambda^{k_1k_2} + y_{i,5}\,\sigma\Lambda\Lambda\,\Lambda^{k_1k_2} + y_{i,6}\,\Lambda^2\left(\eta\sigma\Lambda\Lambda^{k_1}\right)^{k_2} \\
&\quad + y_{i,7}\,\Lambda^2\left(\eta\sigma\Lambda\Lambda^{k_2}\right)^{k_1} + y_{i,8}\,\sigma\Lambda\Lambda\left(\eta\sigma\Lambda\Lambda^{k_1}\right)^{k_2} + y_{i,9}\,\sigma\Lambda\Lambda\left(\eta\sigma\Lambda\Lambda^{k_2}\right)^{k_1} \\
&\quad + y_{i,10}\,\left(\eta\sigma\Lambda\Lambda^{k_3}\right)^{k_1}\Lambda^{k_2}{}_{k_3} + y_{i,11}\,\left(\eta\sigma\Lambda\Lambda^{k_3}\right)^{k_2}\Lambda^{k_1}{}_{k_3}, \qquad \text{(A23)} \\
Y_i^{k_1k_2k_3k_4} &= y_{i,1}\,\eta^{k_1k_2}\,\eta^{k_3k_4}\,\Lambda^2\,\Lambda^2 + y_{i,2}\,\eta^{k_1k_2}\,\eta^{k_3k_4}\,\Lambda^2\,\sigma\Lambda\Lambda + y_{i,3}\,\eta^{k_1k_2}\,\eta^{k_3k_4}\,\sigma\Lambda\Lambda\,\sigma\Lambda\Lambda \\
&\quad + y_{i,4}\,\eta^{k_1k_2}\,\Lambda^2\,\Lambda^{k_3k_4} + y_{i,5}\,\eta^{k_1k_2}\,\sigma\Lambda\Lambda\,\Lambda^{k_3k_4} + y_{i,6}\,\eta^{k_1k_2}\,\Lambda^2\left(\eta\sigma\Lambda\Lambda^{k_3}\right)^{k_4} \\
&\quad + y_{i,7}\,\eta^{k_1k_2}\,\sigma\Lambda\Lambda\,\left(\eta\sigma\Lambda\Lambda^{k_3}\right)^{k_4}, \qquad \text{(A24)} \\
\overline{Y}_i^{k_1,k_2k_3k_4} &= y_{i,1}\,\eta^{k_1k_2}\,\eta^{k_3k_4}\,\Lambda^2\,\Lambda^2 + y_{i,2}\,\eta^{k_1k_2}\,\eta^{k_3k_4}\,\Lambda^2\,\sigma\Lambda\Lambda + y_{i,3}\,\eta^{k_1k_2}\,\eta^{k_3k_4}\,\sigma\Lambda\Lambda\,\sigma\Lambda\Lambda \\
&\quad + y_{i,4}\,\eta^{k_3k_4}\,\Lambda^2\,\Lambda^{k_1k_2} + y_{i,5}\,\eta^{k_1k_2}\,\Lambda^2\,\Lambda^{k_3k_4} + y_{i,6}\,\eta^{k_2k_3}\,\sigma\Lambda\Lambda\,\Lambda^{k_1k_4} \\
&\quad + y_{i,7}\,\eta^{k_1k_2}\,\sigma\Lambda\Lambda\,\Lambda^{k_3k_4} + y_{i,8}\,\eta^{k_2k_3}\,\Lambda^2\left(\eta\sigma\Lambda\Lambda^{k_1}\right)^{k_4} + y_{i,9}\,\eta^{k_2k_3}\,\Lambda^2\left(\eta\sigma\Lambda\Lambda^{k_4}\right)^{k_1} \\
&\quad + y_{i,10}\,\eta^{k_1k_2}\,\Lambda^2\left(\eta\sigma\Lambda\Lambda^{k_3}\right)^{k_4} + y_{i,11}\,\eta^{k_2k_3}\,\sigma\Lambda\Lambda\left(\eta\sigma\Lambda\Lambda^{k_1}\right)^{k_4} \\
&\quad + y_{i,12}\,\eta^{k_2k_3}\,\sigma\Lambda\Lambda\left(\eta\sigma\Lambda\Lambda^{k_4}\right)^{k_1} + y_{i,13}\,\eta^{k_1k_2}\,\sigma\Lambda\Lambda\left(\eta\sigma\Lambda\Lambda^{k_3}\right)^{k_4} \\
&\quad + y_{i,14}\,\Lambda^{k_2k_3}\left(\eta\sigma\Lambda\Lambda^{k_1}\right)^{k_4} + y_{i,15}\,\Lambda^{k_2k_3}\left(\eta\sigma\Lambda\Lambda^{k_4}\right)^{k_1} \\
&\quad + y_{i,16}\,\Lambda^{k_1k_2}\left(\eta\sigma\Lambda\Lambda^{k_3}\right)^{k_4} + y_{i,17}\,\eta^{k_2k_3}\,(\eta\sigma\Lambda\Lambda^{n_1})^{k_4}\,\Lambda^{k_1}{}_{n_1} \\
&\quad + y_{i,18}\,\eta^{k_2k_3}\,(\eta\sigma\Lambda\Lambda^{n_1})^{k_1}\,\Lambda^{k_4}{}_{n_1}, \qquad \text{(A25)} \\
\widetilde{Y}_i^{k_1k_2,k_3k_4} &= y_{i,1}\,\eta^{k_1k_3}\,\eta^{k_2k_4}\,\Lambda^2\Lambda^2 + y_{i,2}\,\eta^{k_1k_2}\,\eta^{k_3k_4}\,\Lambda^2\,\Lambda^2 + y_{i,3}\,\eta^{k_1k_3}\,\eta^{k_2k_4}\,\Lambda^2\,\sigma\Lambda\Lambda \\
&\quad + y_{i,4}\,\eta^{k_1k_2}\,\eta^{k_3k_4}\,\Lambda^2\,\sigma\Lambda\Lambda + y_{i,5}\,\eta^{k_1k_3}\,\eta^{k_2k_4}\,\sigma\Lambda\Lambda\,\sigma\Lambda\Lambda \\
&\quad + y_{i,6}\,\eta^{k_1k_2}\,\eta^{k_3k_4}\,\sigma\Lambda\Lambda\,\sigma\Lambda\Lambda + y_{i,7}\,\eta^{k_1k_3}\,\Lambda^2\,\Lambda^{k_2k_4} + y_{i,8}\,\eta^{k_1k_2}\,\Lambda^2\,\Lambda^{k_3k_4} \\
&\quad + y_{i,9}\,\eta^{k_1k_3}\,\sigma\Lambda\Lambda\,\Lambda^{k_2k_4} + y_{i,10}\,\eta^{k_1k_2}\,\sigma\Lambda\Lambda\,\Lambda^{k_3k_4} + y_{i,11}\,\eta^{k_1k_3}\,\Lambda^2\left(\eta\sigma\Lambda\Lambda^{k_4}\right)^{k_2} \\
&\quad + y_{i,12}\,\eta^{k_1k_2}\,\Lambda^2\left(\eta\sigma\Lambda\Lambda^{k_3}\right)^{k_4} + y_{i,13}\,\eta^{k_1k_3}\,\sigma\Lambda\Lambda\left(\eta\sigma\Lambda\Lambda^{k_4}\right)^{k_2} \\
&\quad + y_{i,14}\,\eta^{k_1k_2}\,\sigma\Lambda\Lambda\left(\eta\sigma\Lambda\Lambda^{k_3}\right)^{k_4} + y_{i,15}\,\Lambda^2\left(\sigma\Lambda\Lambda^{k_1k_3}\right)^{k_2k_4} \\
&\quad + y_{i,16}\,\sigma\Lambda\Lambda\left(\sigma\Lambda\Lambda^{k_1k_3}\right)^{k_2k_4} + y_{i,17}\,\Lambda^{k_1k_2}\,\Lambda^{k_3k_4} + y_{i,18}\,\Lambda^{k_1k_2}\left(\eta\sigma\Lambda\Lambda^{k_3}\right)^{k_4} \\
&\quad + y_{i,19}\,\eta^{k_1k_3}\,(\eta\sigma\Lambda\Lambda^{n_1})^{k_4}\,\Lambda^{k_2}{}_{n_1} + y_{i,20}\,\epsilon^{k_1k_3n_1n_2}\,\Lambda^{k_2}{}_{n_1}\,\Lambda^{k_4}{}_{n_2}, \qquad \text{(A26)} \\
\hat{Y}_i^{k_1k_2,k_3k_4} &= y_{i,1}\,\eta^{k_1k_3}\,\eta^{k_2k_4}\,\Lambda^2\,\Lambda^2 + y_{i,2}\,\eta^{k_1k_2}\,\eta^{k_3k_4}\,\Lambda^2\,\Lambda^2 + y_{i,3}\,\eta^{k_1k_3}\,\eta^{k_2k_4}\,\Lambda^2\,\sigma\Lambda\Lambda \\
&\quad + y_{i,4}\,\eta^{k_1k_2}\eta^{k_3k_4}\,\Lambda^2\,\sigma\Lambda\Lambda + y_{i,5}\,\eta^{k_1k_3}\,\eta^{k_2k_4}\,\sigma\Lambda\Lambda\,\sigma\Lambda\Lambda \\
&\quad + y_{i,6}\,\eta^{k_1k_2}\,\eta^{k_3k_4}\,\sigma\Lambda\Lambda\,\sigma\Lambda\Lambda + y_{i,7}\,\eta^{k_3k_4}\,\Lambda^2\,\Lambda^{k_1k_2} + y_{i,8}\,\eta^{k_1k_3}\,\Lambda^2\,\Lambda^{k_2k_4} \\
&\quad + y_{i,9}\,\eta^{k_1k_2}\,\Lambda^2\,\Lambda^{k_3k_4} + y_{i,10}\,\eta^{k_3k_4}\,\sigma\Lambda\Lambda\,\Lambda^{k_1k_2} + y_{i,11}\,\eta^{k_1k_3}\,\sigma\Lambda\Lambda\,\Lambda^{k_2k_4}
\end{aligned}
$$

$$
\begin{aligned}
&+y_{i,12}\ \eta^{k_1k_2}\ \sigma\Lambda\Lambda\ \Lambda^{k_3k_4} + y_{i,13}\ \eta^{k_1k_3}\ \Lambda^2\left(\eta\sigma\Lambda\Lambda^{k_2}\right)^{k_4}\\
&+y_{i,14}\ \eta^{k_3k_4}\ \Lambda^2\left(\eta\sigma\Lambda\Lambda^{k_1}\right)^{k_2} + y_{i,15}\ \eta^{k_1k_3}\ \Lambda^2\left(\eta\sigma\Lambda\Lambda^{k_4}\right)^{k_2}\\
&+y_{i,16}\ \eta^{k_1k_2}\ \Lambda^2\left(\eta\sigma\Lambda\Lambda^{k_3}\right)^{k_4} + y_{i,17}\ \eta^{k_1k_3}\ \sigma\Lambda\Lambda\left(\eta\sigma\Lambda\Lambda^{k_2}\right)^{k_4}\\
&+y_{i,18}\ \eta^{k_3k_4}\ \sigma\Lambda\Lambda\left(\eta\sigma\Lambda\Lambda^{k_1}\right)^{k_2} + y_{i,19}\ \eta^{k_1k_3}\ \sigma\Lambda\Lambda\left(\eta\sigma\Lambda\Lambda^{k_4}\right)^{k_2}\\
&+y_{i,20}\ \eta^{k_1k_2}\ \sigma\Lambda\Lambda\left(\eta\sigma\Lambda\Lambda^{k_3}\right)^{k_4} + y_{i,21}\ \Lambda^2\left(\sigma\Lambda\Lambda^{k_1k_3}\right)^{k_2k_4}\\
&+y_{i,22}\ \sigma\Lambda\Lambda\left(\sigma\Lambda\Lambda^{k_1k_3}\right)^{k_2k_4} + y_{i,23}\ \Lambda^{k_1k_2}\ \Lambda^{k_3k_4} + y_{i,24}\ \Lambda^{k_3k_4}\left(\eta\sigma\Lambda\Lambda^{k_1}\right)^{k_2}\\
&+y_{i,25}\ \Lambda^{k_1k_2}\left(\eta\sigma\Lambda\Lambda^{k_3}\right)^{k_4} + y_{i,26}\ \eta^{k_1k_3}\left(\eta\sigma\Lambda\Lambda^{n_1}\right)^{k_4}\Lambda^{k_2}{}_{n_1}\\
&+y_{i,27}\ \eta^{k_1k_3}\left(\eta\sigma\Lambda\Lambda^{n_1}\right)^{k_2}\Lambda^{k_4}{}_{n_1} + y_{i,28}\ \epsilon^{k_1k_3n_1n_2}\ \Lambda^{k_2}{}_{n_1}\ \Lambda^{k_4}{}_{n_2}, \qquad \text{(A27)}
\end{aligned}
$$

$$
\begin{aligned}
\widetilde{\overline{Y}}_i{}^{k_1,k_2k_3,k_4} \;=\;& y_{i,1}\ \eta^{k_1k_4}\ \eta^{k_2k_3}\ \Lambda^2\ \Lambda^2 + y_{i,2}\ \eta^{k_1k_3}\ \eta^{k_2k_4}\ \Lambda^2\ \Lambda^2 + y_{i,3}\ \eta^{k_1k_4}\ \eta^{k_2k_3}\ \Lambda^2\ \sigma\Lambda\Lambda\\
&+y_{i,4}\ \eta^{k_1k_3}\ \eta^{k_2k_4}\ \Lambda^2\ \sigma\Lambda\Lambda + y_{i,5}\ \eta^{k_1k_4}\ \eta^{k_2k_3}\ \sigma\Lambda\Lambda\ \sigma\Lambda\Lambda\\
&+y_{i,6}\ \eta^{k_1k_3}\ \eta^{k_2k_4}\ \sigma\Lambda\Lambda\ \sigma\Lambda\Lambda + y_{i,7}\ \eta^{k_2k_3}\ \Lambda^2\ \Lambda^{k_1k_4} + y_{i,8}\ \eta^{k_2k_4}\ \Lambda^2\ \Lambda^{k_1k_3}\\
&+y_{i,9}\ \eta^{k_1k_2}\ \Lambda^2\ \Lambda^{k_3k_4} + y_{i,10}\ \eta^{k_1k_4}\ \Lambda^2\ \Lambda^{k_2k_3} + y_{i,11}\ \eta^{k_2k_3}\ \sigma\Lambda\Lambda\ \Lambda^{k_1k_4}\\
&+y_{i,12}\ \eta^{k_2k_4}\ \sigma\Lambda\Lambda\ \Lambda^{k_1k_3} + y_{i,13}\ \eta^{k_1k_2}\ \sigma\Lambda\Lambda\ \Lambda^{k_3k_4} + y_{i,14}\ \eta^{k_1k_4}\ \sigma\Lambda\Lambda\ \Lambda^{k_2k_3}\\
&+y_{i,15}\ \eta^{k_2k_3}\ \Lambda^2\left(\eta\sigma\Lambda\Lambda^{k_1}\right)^{k_4} + y_{i,16}\ \eta^{k_2k_3}\ \Lambda^2\left(\eta\sigma\Lambda\Lambda^{k_4}\right)^{k_1}\\
&+y_{i,17}\ \eta^{k_2k_4}\ \Lambda^2\left(\eta\sigma\Lambda\Lambda^{k_1}\right)^{k_3} + y_{i,18}\ \eta^{k_1k_2}\ \Lambda^2\left(\eta\sigma\Lambda\Lambda^{k_4}\right)^{k_3}\\
&+y_{i,19}\ \eta^{k_2k_4}\ \Lambda^2\left(\eta\sigma\Lambda\Lambda^{k_3}\right)^{k_1} + y_{i,20}\ \eta^{k_1k_2}\ \Lambda^2\left(\eta\sigma\Lambda\Lambda^{k_3}\right)^{k_4}\\
&+y_{i,21}\ \eta^{k_1k_4}\ \Lambda^2\left(\eta\sigma\Lambda\Lambda^{k_2}\right)^{k_3} + y_{i,22}\ \eta^{k_2k_3}\ \sigma\Lambda\Lambda\left(\eta\sigma\Lambda\Lambda^{k_1}\right)^{k_4}\\
&+y_{i,23}\ \eta^{k_2k_3}\ \sigma\Lambda\Lambda\left(\eta\sigma\Lambda\Lambda^{k_4}\right)^{k_1} + y_{i,24}\ \eta^{k_2k_4}\ \sigma\Lambda\Lambda\left(\eta\sigma\Lambda\Lambda^{k_1}\right)^{k_3}\\
&+y_{i,25}\ \eta^{k_1k_2}\ \sigma\Lambda\Lambda\left(\eta\sigma\Lambda\Lambda^{k_4}\right)^{k_3} + y_{i,26}\ \eta^{k_2k_4}\ \sigma\Lambda\Lambda\left(\eta\sigma\Lambda\Lambda^{k_3}\right)^{k_1}\\
&+y_{i,27}\ \eta^{k_1k_2}\ \sigma\Lambda\Lambda\left(\eta\sigma\Lambda\Lambda^{k_3}\right)^{k_4} + y_{i,28}\ \eta^{k_1k_4}\ \sigma\Lambda\Lambda\left(\eta\sigma\Lambda\Lambda^{k_2}\right)^{k_3}\\
&+y_{i,29}\ \epsilon^{k_1k_2k_4n_1}\ \Lambda^2\ \Lambda^{k_3}{}_{n_1} + y_{i,30}\ \Lambda^2\left(\sigma\Lambda\Lambda^{k_1k_2}\right)^{k_3k_4} + y_{i,31}\ \Lambda^2\left(\sigma\Lambda\Lambda^{k_2k_4}\right)^{k_1k_3}\\
&+y_{i,32}\ \sigma\Lambda\Lambda\left(\sigma\Lambda\Lambda^{k_1k_2}\right)^{k_3k_4} + y_{i,33}\ \sigma\Lambda\Lambda\left(\sigma\Lambda\Lambda^{k_2k_4}\right)^{k_1k_3} + y_{i,34}\ \Lambda^{k_1k_4}\ \Lambda^{k_2k_3}\\
&+y_{i,35}\ \Lambda^{k_2k_3}\left(\eta\sigma\Lambda\Lambda^{k_1}\right)^{k_4} + y_{i,36}\ \Lambda^{k_2k_3}\left(\eta\sigma\Lambda\Lambda^{k_4}\right)^{k_1} + y_{i,37}\ \Lambda^{k_2k_4}\left(\eta\sigma\Lambda\Lambda^{k_3}\right)^{k_1}\\
&+y_{i,38}\ \Lambda^{k_1k_2}\left(\eta\sigma\Lambda\Lambda^{k_3}\right)^{k_4} + y_{i,39}\ \Lambda^{k_1k_4}\left(\eta\sigma\Lambda\Lambda^{k_2}\right)^{k_3}\\
&+y_{i,40}\ \left(\sigma\Lambda\Lambda^{k_2n_1}\right)^{k_3k_4}\Lambda^{k_1}{}_{n_1} + y_{i,41}\ \left(\sigma\Lambda\Lambda^{k_2n_1}\right)^{k_1k_3}\Lambda^{k_4}{}_{n_1}\\
&+y_{i,42}\ \left(\sigma\Lambda\Lambda^{k_2n_1}\right)^{k_1k_4}\Lambda^{k_3}{}_{n_1} + y_{i,43}\ \eta^{k_2k_3}\left(\eta\sigma\Lambda\Lambda^{n_1}\right)^{k_4}\Lambda^{k_1}{}_{n_1}\\
&+y_{i,44}\ \eta^{k_2k_3}\left(\eta\sigma\Lambda\Lambda^{n_1}\right)^{k_1}\Lambda^{k_4}{}_{n_1} + y_{i,45}\ \eta^{k_2k_4}\left(\eta\sigma\Lambda\Lambda^{n_1}\right)^{k_3}\Lambda^{k_1}{}_{n_1}\\
&+y_{i,46}\ \eta^{k_1k_2}\left(\eta\sigma\Lambda\Lambda^{n_1}\right)^{k_3}\Lambda^{k_4}{}_{n_1} + y_{i,47}\ \eta^{k_2k_4}\left(\eta\sigma\Lambda\Lambda^{n_1}\right)^{k_1}\Lambda^{k_3}{}_{n_1}\\
&+y_{i,48}\ \eta^{k_1k_2}\left(\eta\sigma\Lambda\Lambda^{n_1}\right)^{k_4}\Lambda^{k_3}{}_{n_1} + y_{i,49}\ \epsilon^{k_2k_4n_1n_2}\ \Lambda^{k_1}{}_{n_2}\ \Lambda^{k_3}{}_{n_1}
\end{aligned}
$$

$$+y_{i,50}\ \epsilon^{k_1k_2n_1n_2}\ \Lambda^{k_3}{}_{n_2}\ \Lambda^{k_4}{}_{n_1}, \tag{A28}$$

$$\begin{aligned}
Y_i^{k_1\dot\alpha\beta} \;=\;& y_{i,1}\ \Lambda^2\ \Lambda^2\left(\overline\sigma^{k_1}\right)^{\dot\alpha\beta} + y_{i,2}\ \Lambda^2\ \sigma\Lambda\Lambda\left(\overline\sigma^{k_1}\right)^{\dot\alpha\beta} + y_{i,3}\ \sigma\Lambda\Lambda\ \sigma\Lambda\Lambda\left(\overline\sigma^{k_1}\right)^{\dot\alpha\beta}\\
&+y_{i,4}\ \Lambda^2\left(\overline\sigma^{k_2}\right)^{\dot\alpha\beta}\Lambda^{k_1}{}_{k_2} + y_{i,5}\ \sigma\Lambda\Lambda\left(\overline\sigma^{k_2}\right)^{\dot\alpha\beta}\Lambda^{k_1}{}_{k_2}\\
&+y_{i,6}\ \Lambda^2\ \eta_{k_2k_3}\left(\overline\sigma^{k_3}\right)^{\dot\alpha\beta}\left(\eta\sigma\Lambda\Lambda^{k_1}\right)^{k_2} + y_{i,7}\ \Lambda^2\ \eta_{k_2k_3}\left(\overline\sigma^{k_3}\right)^{\dot\alpha\beta}\left(\eta\sigma\Lambda\Lambda^{k_2}\right)^{k_1}\\
&+y_{i,8}\ \sigma\Lambda\Lambda\ \eta_{k_2k_3}\left(\overline\sigma^{k_3}\right)^{\dot\alpha\beta}\left(\eta\sigma\Lambda\Lambda^{k_1}\right)^{k_2} + y_{i,9}\ \sigma\Lambda\Lambda\ \eta_{k_2k_3}\left(\overline\sigma^{k_3}\right)^{\dot\alpha\beta}\left(\eta\sigma\Lambda\Lambda^{k_2}\right)^{k_1}\\
&+y_{i,10}\ \eta_{k_3k_4}\left(\overline\sigma^{k_4}\right)^{\dot\alpha\beta}\left(\eta\sigma\Lambda\Lambda^{k_2}\right)^{k_3}\Lambda^{k_1}{}_{k_2} + y_{i,11}\ \Lambda_{k_2k_3}\left(\overline\sigma^{k_3}\right)^{\dot\alpha\beta}\left(\eta\sigma\Lambda\Lambda^{k_2}\right)^{k_1},
\end{aligned} \tag{A29}$$

$$\begin{aligned}
Y_i^{k_1k_2\alpha\beta} \;=\;& y_{i,1}\ \eta^{k_1k_2}\ \epsilon^{\alpha\beta}\ \Lambda^2\Lambda^2 + y_{i,2}\ \eta^{k_1k_2}\ \epsilon^{\alpha\beta}\ \Lambda^2\ \sigma\Lambda\Lambda + y_{i,3}\ \eta^{k_1k_2}\ \epsilon^{\alpha\beta}\ \sigma\Lambda\Lambda\ \sigma\Lambda\Lambda\\
&+y_{i,4}\ \epsilon^{\alpha\beta}\ \Lambda^2\left(\eta\sigma\Lambda\Lambda^{k_2}\right)^{k_1} + y_{i,5}\ \epsilon^{\alpha\beta}\ \sigma\Lambda\Lambda\left(\eta\sigma\Lambda\Lambda^{k_2}\right)^{k_1} + y_{i,6}\ \epsilon^{\alpha\beta}\ \Lambda^2\ \Lambda^{k_1k_2}\\
&+y_{i,7}\ \epsilon^{\alpha\beta}\ \sigma\Lambda\Lambda\ \Lambda^{k_1k_2} + y_{i,8}\ \Lambda^2\left(\sigma^{k_1k_3}\right)^{\alpha\beta}\Lambda^{k_2}{}_{k_3} + y_{i,9}\ \sigma\Lambda\Lambda\left(\sigma^{k_1k_3}\right)^{\alpha\beta}\Lambda^{k_2}{}_{k_3}\\
&+y_{i,10}\ \Lambda^2\left(\sigma\Lambda^{k_2k_3}\right)^{k_1\alpha}\Lambda_{k_3}{}^{\beta} + y_{i,11}\ \sigma\Lambda\Lambda\left(\sigma\Lambda^{k_2k_3}\right)^{k_1\alpha}\Lambda_{k_3}{}^{\beta}\\
&+y_{i,12}\ \Lambda^2\left(\eta\sigma\Lambda^{k_2}\right)^{\alpha}\Lambda^{k_1\beta} + y_{i,13}\ \sigma\Lambda\Lambda\left(\eta\sigma\Lambda^{k_2}\right)^{\alpha}\Lambda^{k_1\beta}\\
&+y_{i,14}\ \eta^{k_1k_2}\ \Lambda^2\left(\eta\sigma\Lambda^{k_3}\right)^{\alpha}\Lambda_{k_3}{}^{\beta} + y_{i,15}\ \eta^{k_1k_2}\ \sigma\Lambda\Lambda\left(\eta\sigma\Lambda^{k_3}\right)^{\alpha}\Lambda_{k_3}{}^{\beta}\\
&+y_{i,16}\ \Lambda^{k_1k_2}\left(\eta\sigma\Lambda^{k_3}\right)^{\alpha}\Lambda_{k_3}{}^{\beta} + y_{i,17}\ \epsilon^{k_2k_3k_4n_1}\ \Lambda^{k_1}{}_{n_1}\ \Lambda_{k_3}{}^{\beta}\ \Lambda_{k_4}{}^{\alpha},
\end{aligned} \tag{A30}$$

$$\begin{aligned}
\overline{Y}_i{}^{k_1k_2\alpha\beta} \;=\;& y_{i,1}\ \eta^{k_1k_2}\ \epsilon^{\alpha\beta}\ \Lambda^2\ \Lambda^2 + y_{i,2}\ \eta^{k_1k_2}\ \epsilon^{\alpha\beta}\ \Lambda^2\ \sigma\Lambda\Lambda + y_{i,3}\ \eta^{k_1k_2}\ \epsilon^{\alpha\beta}\ \sigma\Lambda\Lambda\ \sigma\Lambda\Lambda\\
&+y_{i,4}\ \Lambda^2\ \Lambda^2\left(\sigma^{k_1k_2}\right)^{\alpha\beta} + y_{i,5}\ \Lambda^2\ \sigma\Lambda\Lambda\left(\sigma^{k_1k_2}\right)^{\alpha\beta} + y_{i,6}\ \sigma\Lambda\Lambda\ \sigma\Lambda\Lambda\left(\sigma^{k_1k_2}\right)^{\alpha\beta}\\
&+y_{i,7}\ \epsilon^{\alpha\beta}\ \Lambda^2\left(\eta\sigma\Lambda\Lambda^{k_2}\right)^{k_1} + y_{i,8}\ \epsilon^{\alpha\beta}\ \sigma\Lambda\Lambda\left(\eta\sigma\Lambda\Lambda^{k_2}\right)^{k_1} + y_{i,9}\ \epsilon^{\alpha\beta}\Lambda^2\Lambda^{k_1k_2}\\
&+y_{i,10}\ \epsilon^{\alpha\beta}\ \sigma\Lambda\Lambda\ \Lambda^{k_1k_2} + y_{i,11}\ \Lambda^2\ \Lambda^{k_1\beta}\ \Lambda^{k_2\alpha} + y_{i,12}\ \sigma\Lambda\Lambda\ \Lambda^{k_1\beta}\ \Lambda^{k_2\alpha}\\
&+y_{i,13}\ \epsilon^{k_1k_2k_3k_4}\ \Lambda^2\ \Lambda_{k_3}{}^{\beta}\ \Lambda_{k_4}{}^{\alpha} + y_{i,14}\ \Lambda^2\left(\sigma^{k_2k_3}\right)^{\alpha\beta}\Lambda^{k_1}{}_{k_3}\\
&+y_{i,15}\ \sigma\Lambda\Lambda\left(\sigma^{k_2k_3}\right)^{\alpha\beta}\Lambda^{k_1}{}_{k_3} + y_{i,16}\ \Lambda^2\left(\sigma\Lambda^{k_2k_3}\right)^{k_1\alpha}\Lambda_{k_3}{}^{\beta}\\
&+y_{i,17}\ \sigma\Lambda\Lambda\left(\sigma\Lambda^{k_2k_3}\right)^{k_1\alpha}\Lambda_{k_3}{}^{\beta} + y_{i,18}\ \Lambda^2\left(\eta\sigma\Lambda^{k_2}\right)^{\alpha}\Lambda^{k_1\beta}\\
&+y_{i,19}\ \sigma\Lambda\Lambda\left(\eta\sigma\Lambda^{k_2}\right)^{\alpha}\Lambda^{k_1\beta} + y_{i,20}\ \epsilon^{\alpha\beta}\left(\eta\sigma\Lambda\Lambda^{k_3}\right)^{k_2}\Lambda^{k_1}{}_{k_3}\\
&+y_{i,21}\ \left(\sigma^{k_3k_4}\right)^{\alpha\beta}\Lambda^{k_1}{}_{k_3}\ \Lambda^{k_2}{}_{k_4} + y_{i,22}\ \epsilon^{k_2k_3k_4n_1}\ \Lambda^{k_1}{}_{n_1}\ \Lambda_{k_3}{}^{\beta}\ \Lambda_{k_4}{}^{\alpha}\\
&+y_{i,23}\ \left(\sigma\Lambda\Lambda^{k_3k_4}\right)^{k_1k_2}\Lambda_{k_3}{}^{\beta}\Lambda_{k_4}{}^{\alpha} + y_{i,24}\ \left(\eta\sigma\Lambda\Lambda^{k_3}\right)^{k_2}\Lambda^{k_1\alpha}\ \Lambda_{k_3}{}^{\beta}\\
&+y_{i,25}\ \left(\sigma\Lambda^{k_3k_4}\right)^{k_2\alpha}\Lambda^{k_1}{}_{k_4}\ \Lambda_{k_3}{}^{\beta} + y_{i,26}\ \left(\eta\sigma\Lambda^{k_3}\right)^{\alpha}\Lambda^{k_2\beta}\ \Lambda^{k_1}{}_{k_3},
\end{aligned} \tag{A31}$$

$$\begin{aligned}
\widetilde{Y}_i^{k_1k_2\alpha\beta} \;=\;& y_{i,1}\ \eta^{k_1k_2}\ \epsilon^{\alpha\beta}\ \Lambda^2\Lambda^2 + y_{i,2}\ \eta^{k_1k_2}\ \epsilon^{\alpha\beta}\ \Lambda^2\ \sigma\Lambda\Lambda + y_{i,3}\ \eta^{k_1k_2}\ \epsilon^{\alpha\beta}\ \sigma\Lambda\Lambda\ \sigma\Lambda\Lambda\\
&+y_{i,4}\ \Lambda^2\ \Lambda^2\left(\sigma^{k_1k_2}\right)^{\alpha\beta} + y_{i,5}\ \Lambda^2\ \sigma\Lambda\Lambda\left(\sigma^{k_1k_2}\right)^{\alpha\beta}\\
&+y_{i,6}\ \sigma\Lambda\Lambda\ \sigma\Lambda\Lambda\left(\sigma^{k_1k_2}\right)^{\alpha\beta} + y_{i,7}\ \epsilon^{\alpha\beta}\ \Lambda^2\left(\eta\sigma\Lambda\Lambda^{k_1}\right)^{k_2}\\
&+y_{i,8}\ \epsilon^{\alpha\beta}\ \Lambda^2\left(\eta\sigma\Lambda\Lambda^{k_2}\right)^{k_1} + y_{i,9}\ \epsilon^{\alpha\beta}\ \sigma\Lambda\Lambda\left(\eta\sigma\Lambda\Lambda^{k_1}\right)^{k_2}\\
&+y_{i,10}\ \epsilon^{\alpha\beta}\ \sigma\Lambda\Lambda\left(\eta\sigma\Lambda\Lambda^{k_2}\right)^{k_1} + y_{i,11}\ \epsilon^{\alpha\beta}\ \Lambda^2\ \Lambda^{k_1k_2} + y_{i,12}\ \epsilon^{\alpha\beta}\ \sigma\Lambda\Lambda\ \Lambda^{k_1k_2}
\end{aligned}$$

$$
\begin{aligned}
&+y_{i,13}\ \Lambda^2\ \Lambda^{k_1\alpha}\ \Lambda^{k_2\beta}+y_{i,14}\ \sigma\Lambda\Lambda\ \Lambda^{k_1\alpha}\ \Lambda^{k_2\beta}+y_{i,15}\ \epsilon^{k_1k_2k_3k_4}\ \Lambda^2\ \Lambda_{k_3}{}^{\beta}\ \Lambda_{k_4}{}^{\alpha}\\
&+y_{i,16}\ \Lambda^2\left(\sigma^{k_1k_3}\right)^{\alpha\beta}\Lambda^{k_2}{}_{k_3}+y_{i,17}\ \Lambda^2\left(\sigma^{k_2k_3}\right)^{\alpha\beta}\Lambda^{k_1}{}_{k_3}\\
&+y_{i,18}\ \sigma\Lambda\Lambda\left(\sigma^{k_1k_3}\right)^{\alpha\beta}\Lambda^{k_2}{}_{k_3}+y_{i,19}\ \sigma\Lambda\Lambda\left(\sigma^{k_2k_3}\right)^{\alpha\beta}\Lambda^{k_1}{}_{k_3}\\
&+y_{i,20}\ \Lambda^2\left(\sigma\Lambda^{k_1k_3}\right)^{k_2\alpha}\Lambda_{k_3}{}^{\beta}+y_{i,21}\ \Lambda^2\left(\sigma\Lambda^{k_2k_3}\right)^{k_1\alpha}\Lambda_{k_3}{}^{\beta}\\
&+y_{i,22}\ \sigma\Lambda\Lambda\left(\sigma\Lambda^{k_1k_3}\right)^{k_2\alpha}\Lambda_{k_3}{}^{\beta}+y_{i,23}\ \sigma\Lambda\Lambda\left(\sigma\Lambda^{k_2k_3}\right)^{k_1\alpha}\Lambda_{k_3}{}^{\beta}\\
&+y_{i,24}\ \Lambda^2\left(\eta\sigma\Lambda^{k_1}\right)^{\alpha}\Lambda^{k_2\beta}+y_{i,25}\ \Lambda^2\left(\eta\sigma\Lambda^{k_2}\right)^{\alpha}\Lambda^{k_1\beta}\\
&+y_{i,26}\ \sigma\Lambda\Lambda\left(\eta\sigma\Lambda^{k_1}\right)^{\alpha}\Lambda^{k_2\beta}+y_{i,27}\ \sigma\Lambda\Lambda\left(\eta\sigma\Lambda^{k_2}\right)^{\alpha}\Lambda^{k_1\beta}\\
&+y_{i,28}\ \eta^{k_1k_2}\ \Lambda^2\left(\eta\sigma\Lambda^{k_3}\right)^{\alpha}\Lambda_{k_3}{}^{\beta}+y_{i,29}\ \eta^{k_1k_2}\ \sigma\Lambda\Lambda\left(\eta\sigma\Lambda^{k_3}\right)^{\alpha}\Lambda_{k_3}{}^{\beta}\\
&+y_{i,30}\ \epsilon^{\alpha\beta}\left(\eta\sigma\Lambda\Lambda^{k_3}\right)^{k_1}\Lambda^{k_2}{}_{k_3}+y_{i,31}\ \epsilon^{\alpha\beta}\left(\eta\sigma\Lambda\Lambda^{k_3}\right)^{k_2}\Lambda^{k_1}{}_{k_3}\\
&+y_{i,32}\ \left(\sigma^{k_3k_4}\right)^{\alpha\beta}\Lambda^{k_1}{}_{k_3}\ \Lambda^{k_2}{}_{k_4}+y_{i,33}\ \Lambda^{k_1k_2}\left(\eta\sigma\Lambda^{k_3}\right)^{\alpha}\Lambda_{k_3}{}^{\beta}\\
&+y_{i,34}\ \epsilon^{k_2k_3k_4n_1}\ \Lambda^{k_1}{}_{n_1}\ \Lambda_{k_3}{}^{\beta}\ \Lambda_{k_4}{}^{\alpha}+y_{i,35}\ \epsilon^{k_1k_3k_4n_1}\ \Lambda^{k_2}{}_{n_1}\ \Lambda_{k_3}{}^{\beta}\ \Lambda_{k_4}{}^{\alpha}\\
&+y_{i,36}\ \left(\sigma\Lambda\Lambda^{k_3k_4}\right)^{k_1k_2}\Lambda_{k_3}{}^{\beta}\ \Lambda_{k_4}{}^{\alpha}+y_{i,37}\ \left(\eta\sigma\Lambda\Lambda^{k_3}\right)^{k_1}\Lambda^{k_2\alpha}\ \Lambda_{k_3}{}^{\beta}\\
&+y_{i,38}\ \left(\eta\sigma\Lambda\Lambda^{k_3}\right)^{k_2}\Lambda^{k_1\alpha}\ \Lambda_{k_3}{}^{\beta}+y_{i,39}\ \left(\sigma\Lambda^{k_3k_4}\right)^{k_1\alpha}\Lambda^{k_2}{}_{k_4}\ \Lambda_{k_3}{}^{\beta}\\
&+y_{i,40}\ \left(\sigma\Lambda^{k_3k_4}\right)^{k_2\alpha}\Lambda^{k_1}{}_{k_4}\ \Lambda_{k_3}{}^{\beta}+y_{i,41}\ \left(\eta\sigma\Lambda^{k_3}\right)^{\alpha}\Lambda^{k_1\beta}\Lambda^{k_2}{}_{k_3}\\
&+y_{i,42}\ \left(\eta\sigma\Lambda^{k_3}\right)^{\alpha}\Lambda^{k_2\beta}\Lambda^{k_1}{}_{k_3}. \qquad \text{(A32)}
\end{aligned}
$$

where $y_{i,j}$ are free parameters. They can combine with operators as follows,

$$
\begin{aligned}
&Y_i^{k_1k_2}\ \left(D^2\Phi\right)\Phi\left(\partial_{k_1}\partial_{k_2}\Phi^+\right),\quad \overline{Y}_i{}^{k_1k_2}\ \left(\partial_{k_1}D^2\Phi\right)\Phi\left(\partial_{k_2}\Phi^+\right)\Phi^+\\
&Y_i^{k_1k_2k_3k_4}\ \Phi\ \left(\partial_{k_1}\partial_{k_2}\partial_{k_3}\partial_{k_4}\Phi^+\right)\Phi^+,\quad \overline{Y}_i{}^{k_1,k_2k_3k_4}\ \Phi\left(\partial_{k_2}\partial_{k_3}\partial_{k_4}\Phi^+\right)\left(\partial_{k_1}\Phi^+\right)\\
&\widetilde{Y}_i^{k_1k_2,k_3k_4}\ \Phi\left(\partial_{k_1}\partial_{k_2}\Phi^+\right)\left(\partial_{k_3}\partial_{k_4}\Phi^+\right),\quad \hat{Y}_i^{k_1k_2,k_3k_4}\left(\partial_{k_3}\partial_{k_4}D^2\Phi\right)\Phi\left(\partial_{k_1}\partial_{k_2}\Phi^+\right)\\
&\overline{\widetilde{Y}}_i{}^{k_1,k_2k_3,k_4}\left(\partial_{k_4}D^2\Phi\right)\Phi\left(\partial_{k_2}\partial_{k_3}\Phi^+\right)\left(\partial_{k_1}\Phi^+\right),\quad Y_i^{k_1\dot\alpha\beta}\left(D_\beta\Phi\right)\left(\partial_{k_1}\Phi\right)\left(\overline{D}_{\dot\alpha}\Phi^+\right)\Phi^+\ \Phi^+\\
&Y_i^{k_1k_2\alpha\beta}\left(\partial_{k_1}\partial_{k_2}D_\alpha\Phi\right)\left(D_\beta\Phi\right)\Phi\ \Phi^+,\quad \overline{Y}_i{}^{k_1k_2\alpha\beta}\left(\partial_{k_1}D_\alpha\Phi\right)\left(\partial_{k_2}D_\beta\Phi\right)\Phi\ \Phi^+\\
&\widetilde{Y}_i^{k_1k_2\alpha\beta}\left(\partial_{k_1}D_\alpha\Phi\right)\left(D_\beta\Phi\right)\left(\partial_{k_2}\Phi\right)\Phi^+ \qquad \text{(A33)}
\end{aligned}
$$

By using Eq. (7) we can construct the following action,

$$
\begin{aligned}
&\int d^4x\ d^4\theta\ \Big\{Y_1\ \overline{\theta}^2\left(D^2\Phi\right)\left(D^2\Phi\right)\Phi+Y_2\ \overline{\theta}^2\left(D^2\Phi\right)\Phi\ \left(\Phi^+\right)^2+Y_3\ \overline{\theta}^2\left(D^2\Phi\right)\left(D^2\Phi\right)\Phi\ \Phi^+\\
&+Y_4\ \overline{\theta}^2\left(D^2\Phi\right)\left(D^2\Phi\right)\left(D^2\Phi\right)\Phi+Y_5\ \overline{\theta}^2\ \Phi\ \left(\Phi^+\right)^4+Y_6\ \overline{\theta}^2\left(D^2\Phi\right)\Phi\ \left(\Phi^+\right)^3\\
&+Y_7\ \overline{\theta}^2\left(D^2\Phi\right)\left(D^2\Phi\right)\Phi\ \left(\Phi^+\right)^2+Y_8\ \overline{\theta}^2\ \Phi\ \left(\Phi^+\right)^5+Y_9\ \overline{\theta}^2\left(D^2\Phi\right)\Phi\ \left(\Phi^+\right)^4\\
&+Y_{10}\ \overline{\theta}^2\ \Phi\ \left(\Phi^+\right)^6\Big\}. \qquad \text{(A34)}
\end{aligned}
$$

By using Eq. (8) we obtain,

$$
\int d^4x\ d^4\theta\ \Big\{Y_{11}{}^{k_1k_2}\ \overline{\theta}^2\left(D^2\Phi\right)\Phi\left(\partial_{k_1}\partial_{k_2}\Phi^+\right)+Y_{12}{}^{k_1k_2}\ \overline{\theta}^2\left(\partial_{k_1}\partial_{k_2}D^2\Phi\right)\Phi\ \Phi^+
$$

$$
\begin{aligned}
&+\overline{Y}_{13}{}^{k_1k_2}\ \overline{\theta}^2\left(\partial_{k_1}D^2\Phi\right)\Phi\left(\partial_{k_2}\Phi^+\right)+Y_{14}{}^{k_1k_2}\ \overline{\theta}^2\left(\partial_{k_1}D^2\Phi\right)\left(\partial_{k_2}D^2\Phi\right)\Phi\\
&+Y_{15}{}^{k_1k_2}\ \overline{\theta}^2\left(\partial_{k_1}\partial_{k_2}D^2\Phi\right)\left(D^2\Phi\right)\Phi+Y_{16}{}^{k_1k_2}\ \overline{\theta}^2\ \Phi\left(\partial_{k_1}\Phi^+\right)\left(\partial_{k_2}\Phi^+\right)\Phi^+\\
&+Y_{17}{}^{k_1k_2}\ \overline{\theta}^2\ \Phi\left(\partial_{k_1}\partial_{k_2}\Phi^+\right)\left(\Phi^+\right)^2+Y_{18}{}^{k_1k_2}\ \overline{\theta}^2\left(D^2\Phi\right)\Phi\left(\partial_{k_1}\Phi^+\right)\left(\partial_{k_2}\Phi^+\right)\\
&+Y_{19}{}^{k_1k_2}\ \overline{\theta}^2\left(D^2\Phi\right)\Phi\left(\partial_{k_1}\partial_{k_2}\Phi^+\right)\Phi^++Y_{20}{}^{k_1k_2}\ \overline{\theta}^2\left(\partial_{k_1}\partial_{k_2}D^2\right)\Phi^2\left(\Phi^+\right)^2\\
&+\overline{Y}_{21}{}^{k_1k_2}\ \overline{\theta}^2\left(\partial_{k_1}D^2\Phi\right)\Phi\left(\partial_{k_2}\Phi^+\right)\Phi^++Y_{22}{}^{k_1k_2}\ \overline{\theta}^2\left(D^2\Phi\right)\left(D^2\Phi\right)\Phi\left(\partial_{k_1}\partial_{k_2}\Phi^+\right)\\
&+Y_{23}{}^{k_1k_2}\ \overline{\theta}^2\left(\partial_{k_1}D^2\Phi\right)\left(\partial_{k_2}D^2\Phi\right)\Phi\ \Phi^++Y_{24}{}^{k_1k_2}\ \overline{\theta}^2\left(\partial_{k_1}\partial_{k_2}D^2\Phi\right)\left(D^2\Phi\right)\Phi\ \Phi^+\\
&+\overline{Y}_{25}{}^{k_1k_2}\ \overline{\theta}^2\left(\partial_{k_1}D^2\Phi\right)\left(D^2\Phi\right)\Phi\left(\partial_{k_2}\Phi^+\right)+Y_{26}{}^{k_1k_2}\ \overline{\theta}^2\ \Phi\left(\partial_{k_1}\Phi^+\right)\left(\partial_{k_2}\Phi^+\right)\left(\Phi^+\right)^2\\
&+Y_{27}{}^{k_1k_2}\ \overline{\theta}^2\ \Phi\left(\partial_{k_1}\partial_{k_2}\Phi^+\right)\left(\Phi^+\right)^3+Y_{28}{}^{k_1k_2}\ \theta^4\left(\partial_{k_1}\partial_{k_2}\Phi^+\right)\left(\Phi^+\right)^4\\
&+Y_{29}{}^{k_1k_2}\ \overline{\theta}^2\left(D^2\Phi\right)\Phi\left(\partial_{k_1}\Phi^+\right)\left(\partial_{k_2}\Phi^+\right)\Phi^++Y_{30}{}^{k_1k_2}\ \overline{\theta}^2\left(D^2\Phi\right)\Phi\left(\partial_{k_1}\partial_{k_2}\Phi^+\right)\left(\Phi^+\right)^2\\
&+Y_{31}{}^{k_1k_2}\ \overline{\theta}^2\left(\partial_{k_1}\partial_{k_2}D^2\Phi\right)\Phi\ \left(\Phi^+\right)^3+\overline{Y}_{32}{}^{k_1k_2}\ \overline{\theta}^2\left(\partial_{k_2}D^2\Phi\right)\Phi\left(\partial_{k_1}\Phi^+\right)\left(\Phi^+\right)^2\\
&+Y_{33}{}^{k_1k_2}\ \overline{\theta}^2\ \Phi\left(\partial_{k_1}\Phi^+\right)\left(\partial_{k_2}\Phi^+\right)\left(\Phi^+\right)^3+Y_{34}{}^{k_1k_2}\ \overline{\theta}^2\ \Phi\left(\partial_{k_1}\partial_{k_2}\Phi^+\right)\left(\Phi^+\right)^4\\
&+Y_{35}{}^{k_1k_2}\ \theta^4\left(\partial_{k_1}\partial_{k_2}\Phi^+\right)\left(\Phi^+\right)^5+Y_{36}{}^{k_1k_2}\ \theta^4\left(\partial_{k_1}\partial_{k_2}\Phi^+\right)\left(\Phi^+\right)^6\Bigg\}. \qquad \text{(A35)}
\end{aligned}
$$

By using Eq. (9) we obtain,

$$
\begin{aligned}
&\int d^4x\ d^4\theta\left\{Y_{37}{}^{k_1k_2k_3k_4}\ \overline{\theta}^2\left(\partial_{k_1}\partial_{k_2}\partial_{k_3}\partial_{k_4}D^2\Phi\right)\Phi+Y_{38}{}^{k_1k_2k_3k_4}\ \overline{\theta}^2\ \Phi\left(\partial_{k_1}\partial_{k_2}\partial_{k_3}\partial_{k_4}\Phi^+\right)\Phi^+\right.\\
&+\overline{Y}_{39}{}^{k_1,k_2k_3k_4}\ \overline{\theta}^2\ \Phi\left(\partial_{k_2}\partial_{k_3}\partial_{k_4}\Phi^+\right)\left(\partial_{k_1}\Phi^+\right)+\widetilde{Y}_{40}{}^{k_1k_2,k_3k_4}\ \overline{\theta}^2\ \Phi\left(\partial_{k_1}\partial_{k_2}\Phi^+\right)\left(\partial_{k_3}\partial_{k_4}\Phi^+\right)\\
&+Y_{41}{}^{k_1k_2k_3k_4}\ \overline{\theta}^2\left(\partial_{k_1}\partial_{k_2}\partial_{k_3}\partial_{k_4}D^2\Phi\right)\Phi\ \Phi^++\overline{Y}_{42}{}^{k_1,k_2k_3k_4}\ \overline{\theta}^2\left(\partial_{k_2}\partial_{k_3}\partial_{k_4}D^2\Phi\right)\Phi\left(\partial_{k_1}\Phi^+\right)\\
&+\hat{Y}_{43}{}^{k_1k_2,k_3k_4}\ \overline{\theta}^2\left(\partial_{k_3}\partial_{k_4}D^2\Phi\right)\Phi\left(\partial_{k_1}\partial_{k_2}\Phi^+\right)+\overline{Y}_{44}{}^{k_1,k_2k_3k_4}\ \overline{\theta}^2\left(\partial_{k_1}D^2\Phi\right)\Phi\left(\partial_{k_2}\partial_{k_3}\partial_{k_4}\Phi^+\right)\\
&+Y_{45}{}^{k_1k_2k_3k_4}\ \overline{\theta}^2\left(D^2\Phi\right)\Phi\left(\partial_{k_1}\partial_{k_2}\partial_{k_3}\partial_{k_4}\Phi^+\right)\\
&+Y_{46}{}^{k_1k_2k_3k_4}\ \overline{\theta}^2\left(\partial_{k_1}\partial_{k_2}\partial_{k_3}\partial_{k_4}D^2\Phi\right)\left(D^2\Phi\right)\Phi\\
&+\overline{Y}_{47}{}^{k_1,k_2k_3k_4}\ \overline{\theta}^2\left(\partial_{k_2}\partial_{k_3}\partial_{k_4}D^2\Phi\right)\left(\partial_{k_1}D^2\Phi\right)\Phi\\
&+\widetilde{Y}_{48}{}^{k_1k_2,k_3k_4}\ \overline{\theta}^2\left(\partial_{k_1}\partial_{k_2}D^2\Phi\right)\left(\partial_{k_3}\partial_{k_4}D^2\Phi\right)\Phi\\
&+Y_{49}{}^{k_1k_2k_3k_4}\ \theta^4\left(\partial_{k_1}\partial_{k_2}\partial_{k_3}\partial_{k_4}\Phi^+\right)\left(\Phi^+\right)^3\\
&+\widetilde{Y}_{50}{}^{k_1k_2,k_3k_4}\ \theta^4\left(\partial_{k_1}\partial_{k_2}\Phi^+\right)\left(\partial_{k_3}\partial_{k_4}\Phi^+\right)\left(\Phi^+\right)^2\\
&+Y_{51}{}^{k_1k_2k_3k_4}\ \overline{\theta}^2\ \Phi\left(\partial_{k_1}\partial_{k_2}\partial_{k_3}\partial_{k_4}\Phi^+\right)\left(\Phi^+\right)^2\\
&+\overline{Y}_{52}{}^{k_1,k_2k_3k_4}\ \overline{\theta}^2\ \Phi\left(\partial_{k_2}\partial_{k_3}\partial_{k_4}\Phi^+\right)\left(\partial_{k_1}\Phi^+\right)\Phi^+\\
&+\hat{Y}_{53}{}^{k_1k_2,k_3k_4}\ \overline{\theta}^2\ \Phi\left(\partial_{k_3}\partial_{k_4}\Phi^+\right)\left(\partial_{k_1}\Phi^+\right)\left(\partial_{k_2}\Phi^+\right)\\
&+\widetilde{Y}_{54}{}^{k_1k_2,k_3k_4}\ \overline{\theta}^2\ \Phi\left(\partial_{k_1}\partial_{k_2}\Phi^+\right)\left(\partial_{k_3}\partial_{k_4}\Phi^+\right)\Phi^+\\
&+Y_{55}{}^{k_1k_2k_3k_4}\ \overline{\theta}^2\left(\partial_{k_1}\partial_{k_2}\partial_{k_3}\partial_{k_4}D^2\Phi\right)\Phi\ \left(\Phi^+\right)^2\\
&+\overline{Y}_{56}{}^{k_1,k_2k_3k_4}\ \overline{\theta}^2\left(\partial_{k_2}\partial_{k_3}\partial_{k_4}D^2\Phi\right)\Phi\left(\partial_{k_1}\Phi^+\right)\Phi^+\\
&+\hat{Y}_{57}{}^{k_1k_2,k_3k_4}\ \overline{\theta}^2\left(\partial_{k_3}\partial_{k_4}D^2\Phi\right)\Phi\left(\partial_{k_1}\partial_{k_2}\Phi^+\right)\Phi^+
\end{aligned}
$$

$$
\begin{aligned}
&+\hat{Y}_{58}{}^{k_1k_2,k_3k_4}\ \overline{\theta}^2\left(\partial_{k_3}\partial_{k_4}D^2\Phi\right)\Phi\left(\partial_{k_1}\Phi^+\right)\left(\partial_{k_2}\Phi^+\right)\\
&+\overline{Y}_{59}{}^{k_1,k_2k_3k_4}\ \overline{\theta}^2\left(\partial_{k_1}D^2\Phi\right)\Phi\left(\partial_{k_2}\partial_{k_3}\partial_{k_4}\Phi^+\right)\Phi^+\\
&+\widetilde{\overline{Y}}_{60}{}^{k_1,k_2k_3,k_4}\ \overline{\theta}^2\left(\partial_{k_4}D^2\Phi\right)\Phi\left(\partial_{k_2}\partial_{k_3}\Phi^+\right)\left(\partial_{k_1}\Phi^+\right)\\
&+Y_{61}{}^{k_1k_2k_3k_4}\ \overline{\theta}^2\left(D^2\Phi\right)\Phi\left(\partial_{k_1}\partial_{k_2}\partial_{k_3}\partial_{k_4}\Phi^+\right)\Phi^+\\
&+\overline{Y}_{62}{}^{k_1,k_2k_3k_4}\ \overline{\theta}^2\left(D^2\Phi\right)\Phi\left(\partial_{k_2}\partial_{k_3}\partial_{k_4}\Phi^+\right)\left(\partial_{k_1}\Phi^+\right)\\
&+\widetilde{Y}_{63}{}^{k_1k_2,k_3k_4}\ \overline{\theta}^2\left(D^2\Phi\right)\Phi\left(\partial_{k_1}\partial_{k_2}\Phi^+\right)\left(\partial_{k_3}\partial_{k_4}\Phi^+\right)\\
&+Y_{64}{}^{k_1k_2k_3k_4}\ \theta^4\left(\partial_{k_1}\partial_{k_2}\partial_{k_3}\partial_{k_4}\Phi^+\right)\left(\Phi^+\right)^4\\
&+\widetilde{Y}_{65}{}^{k_1k_2,k_3k_4}\ \theta^4\left(\partial_{k_1}\partial_{k_2}\Phi^+\right)\left(\partial_{k_3}\partial_{k_4}\Phi^+\right)\left(\Phi^+\right)^3\\
&+Y_{66}{}^{k_1k_2k_3k_4}\ \overline{\theta}^2\ \Phi\left(\partial_{k_1}\partial_{k_2}\partial_{k_3}\partial_{k_4}\Phi^+\right)\left(\Phi^+\right)^3\\
&+\overline{Y}_{67}{}^{k_1,k_2k_3k_4}\ \overline{\theta}^2\ \Phi\left(\partial_{k_2}\partial_{k_3}\partial_{k_4}\Phi^+\right)\left(\partial_{k_1}\Phi^+\right)\left(\Phi^+\right)^2\\
&+\widetilde{Y}_{68}{}^{k_1k_2,k_3k_4}\ \overline{\theta}^2\ \Phi\left(\partial_{k_1}\partial_{k_2}\Phi^+\right)\left(\partial_{k_3}\partial_{k_4}\Phi^+\right)\left(\Phi^+\right)^2\\
&+\hat{Y}_{69}{}^{k_1k_2,k_3k_4}\ \overline{\theta}^2\ \Phi\left(\partial_{k_3}\partial_{k_4}\Phi^+\right)\left(\partial_{k_1}\Phi^+\right)\left(\partial_{k_2}\Phi^+\right)\Phi^+\\
&+Y_{70}{}^{k_1k_2k_3k_4}\ \overline{\theta}^2\ \Phi\left(\partial_{k_1}\Phi^+\right)\left(\partial_{k_2}\Phi^+\right)\left(\partial_{k_3}\Phi^+\right)\left(\partial_{k_4}\Phi^+\right)\\
&+Y_{71}{}^{k_1k_2k_3k_4}\ \theta^4\left(\partial_{k_1}\partial_{k_2}\partial_{k_3}\partial_{k_4}\Phi^+\right)\left(\Phi^+\right)^5\\
&+\widetilde{Y}_{72}{}^{k_1k_2,k_3k_4}\ \theta^4\left(\partial_{k_1}\partial_{k_2}\Phi^+\right)\left(\partial_{k_3}\partial_{k_4}\Phi^+\right)\left(\Phi^+\right)^4\Bigg\}.
\end{aligned}
\tag{A36}
$$

By using Eq. (10) we obtain,

$$
\begin{aligned}
&\int d^4x\ d^4\theta\ \Bigg\{Y_{73}\ \overline{\theta}^2\left(D^2\Phi\right)\left(D^2\Phi\right)\Phi^2+Y_{74}\ \overline{\theta}^2\left(D^2\Phi\right)\Phi^2\left(\Phi^+\right)^2+Y_{75}\ \overline{\theta}^2\left(D^2\Phi\right)\left(D^2\Phi\right)\Phi^3\\
&+Y_{76}\ \overline{\theta}^2\left(D^2\Phi\right)\left(D^2\Phi\right)\Phi^2\ \Phi^++Y_{77}\ \overline{\theta}^2\left(D^2\Phi\right)\left(D^2\Phi\right)\left(D^2\Phi\right)\Phi^2+Y_{78}\ \overline{\theta}^2\ \Phi^2\left(\Phi^+\right)^4\\
&+Y_{79}\ \overline{\theta}^2\left(D^2\Phi\right)\Phi^3\left(\Phi^+\right)^2+Y_{80}\ \overline{\theta}^2\left(D^2\Phi\right)\Phi^2\left(\Phi^+\right)^3+Y_{81}\ \overline{\theta}^2\left(D^2\Phi\right)\left(D^2\Phi\right)\Phi^3\ \Phi^+\\
&+Y_{82}\ \overline{\theta}^2\left(D^2\Phi\right)\left(D^2\Phi\right)\Phi^2\left(\Phi^+\right)^2+Y_{83}\ \overline{\theta}^2\left(D^2\Phi\right)\left(D^2\Phi\right)\left(D^2\Phi\right)\Phi^3+Y_{84}\ \overline{\theta}^2\ \Phi^3\left(\Phi^+\right)^4\\
&+Y_{85}\ \overline{\theta}^2\ \Phi^2\left(\Phi^+\right)^5+Y_{86}\ \overline{\theta}^2\left(D^2\Phi\right)\Phi^3\left(\Phi^+\right)^3+Y_{87}\ \overline{\theta}^2\left(D^2\Phi\right)\Phi^2\left(\Phi^+\right)^4\\
&+Y_{88}\ \overline{\theta}^2\left(D^2\Phi\right)\left(D^2\Phi\right)\Phi^3\left(\Phi^+\right)^2+Y_{89}\ \overline{\theta}^2\ \Phi^3\left(\Phi^+\right)^5+Y_{90}\ \overline{\theta}^2\ \Phi^2\left(\Phi^+\right)^6\\
&+Y_{91}\ \overline{\theta}^2\left(D^2\Phi\right)\Phi^3\left(\Phi^+\right)^4+Y_{92}\ \overline{\theta}^2\ \Phi^3\left(\Phi^+\right)^6\Bigg\}.
\end{aligned}
\tag{A37}
$$

By using Eq. (11) we obtain,

$$
\begin{aligned}
&\int d^4x\ d^4\theta\ \Bigg\{Y_{93}{}^{k_1k_2}\ \overline{\theta}^2\ \left(\partial_{k_2}\partial_{k_1}D^2\Phi\right)\left(D^2\Phi\right)\Phi^2+Y_{94}{}^{k_1k_2}\ \overline{\theta}^2\ \left(\partial_{k_1}D^2\Phi\right)\left(\partial_{k_2}D^2\Phi\right)\Phi^2\\
&+Y_{95}{}^{k_1k_2}\ \overline{\theta}^2\left(D^2\Phi\right)\left(D^2\Phi\right)\left(\partial_{k_1}\partial_{k_2}\Phi\right)\Phi+Y_{96}{}^{k_1k_2\alpha\beta}\ \overline{\theta}^2\ \left(\partial_{k_2}\partial_{k_1}D_\alpha\Phi\right)\left(D_\beta\Phi\right)\left(D^2\Phi\right)\Phi\\
&+\overline{Y}_{97}{}^{k_1k_2\alpha\beta}\ \overline{\theta}^2\ \left(\partial_{k_1}D_\alpha\Phi\right)\left(\partial_{k_2}D_\beta\Phi\right)\left(D^2\Phi\right)\Phi+\widetilde{Y}_{98}{}^{k_1k_2\alpha\beta}\ \overline{\theta}^2\ \left(\partial_{k_2}D_\beta\Phi\right)\left(D_\alpha\Phi\right)\left(\partial_{k_1}D^2\Phi\right)\Phi\\
&+Y_{99}{}^{k_1k_2}\ \epsilon^{\alpha\beta}\ \overline{\theta}^2\left(D_\alpha\Phi\right)\left(D_\beta\Phi\right)\left(\partial_{k_2}\partial_{k_1}D^2\Phi\right)\Phi+Y_{100}\ \overline{\theta}^2\left(D^2\Phi\right)\left(D^2\Phi\right)\Phi\left(\overline{D}^2\Phi^+\right)\\
&+Y_{101}\ \epsilon^{\alpha\beta}\ \overline{\theta}^2\left(D_\alpha\Phi\right)\left(D_\beta\Phi\right)\left(D^2\Phi\right)\left(\overline{D}^2\Phi^+\right)+Y_{102}{}^{k_1\dot{\alpha}\beta}\ \overline{\theta}^2\ \left(\partial_{k_1}D_\beta\Phi\right)\left(D^2\Phi\right)\Phi\left(\overline{D}_{\dot{\alpha}}\Phi^+\right)
\end{aligned}
$$

$$
\begin{aligned}
&+Y_{103}{}^{k_1\dot\alpha\beta}\ \overline\theta^2\,(D_\beta\Phi)\left(\partial_{k_1}D^2\Phi\right)\Phi\left(\overline D_{\dot\alpha}\Phi^+\right)+Y_{104}{}^{k_1\dot\alpha\beta}\ \overline\theta^2\,(D_\beta\Phi)\left(D^2\Phi\right)(\partial_{k_1}\Phi)\left(\overline D_{\dot\alpha}\Phi^+\right)\\
&+Y_{105}{}^{k_1k_2}\ \overline\theta^2\ \left(\partial_{k_2}\partial_{k_1}D^2\Phi\right)\Phi^2\ \Phi^++\overline Y_{106}{}^{k_1k_2}\ \overline\theta^2\ \left(\partial_{k_1}D^2\Phi\right)(\partial_{k_2}\Phi)\,\Phi\ \Phi^+\\
&+Y_{107}{}^{k_1k_2}\ \overline\theta^2\left(D^2\Phi\right)(\partial_{k_1}\partial_{k_2}\Phi)\,\Phi\ \Phi^++Y_{108}{}^{k_1k_2}\ \overline\theta^2\left(D^2\Phi\right)(\partial_{k_1}\Phi)\,(\partial_{k_2}\Phi)\,\Phi^+\\
&+Y_{109}{}^{k_1k_2\alpha\beta}\ \overline\theta^2\ \ (\partial_{k_2}\partial_{k_1}D_\alpha\Phi)\,(D_\beta\Phi)\,\Phi\ \Phi^++\overline Y_{110}{}^{k_1k_2\alpha\beta}\ \overline\theta^2\ \ (\partial_{k_1}D_\alpha\Phi)\,(\partial_{k_2}D_\beta\Phi)\,\Phi\ \Phi^+\\
&+\widetilde Y_{111}{}^{k_1k_2\alpha\beta}\ \overline\theta^2\ \ (\partial_{k_2}D_\beta\Phi)\,(D_\alpha\Phi)\,(\partial_{k_1}\Phi)\,\Phi^++Y_{112}{}^{k_1k_2}\ \epsilon^{\alpha\beta}\ \overline\theta^2\,(D_\alpha\Phi)\,(D_\beta\Phi)\,(\partial_{k_1}\partial_{k_2}\Phi)\,\Phi^+\\
&+Y_{113}\ \overline\theta^2\left(D^2\Phi\right)\Phi\left(\overline D^2\Phi^+\right)\left(\Phi^+\right)^2+Y_{114}\ \epsilon^{\alpha\beta}\ \overline\theta^2\,(D_\alpha\Phi)\,(D_\beta\Phi)\left(\overline D^2\Phi^+\right)\left(\Phi^+\right)^2\\
&+Y_{115}\ \epsilon^{\dot\alpha\dot\beta}\ \overline\theta^2\left(D^2\Phi\right)\Phi\left(\overline D_{\dot\alpha}\Phi^+\right)\left(\overline D_{\dot\beta}\Phi^+\right)\Phi^++Y_{116}{}^{k_1\dot\alpha\beta}\ \overline\theta^2\ \ (\partial_{k_1}D_\beta\Phi)\,\Phi\left(\overline D_{\dot\alpha}\Phi^+\right)\left(\Phi^+\right)^2\\
&+Y_{117}{}^{k_1\dot\alpha\beta}\ \overline\theta^2\,(D_\beta\Phi)\,(\partial_{k_1}\Phi)\left(\overline D_{\dot\alpha}\Phi^+\right)\left(\Phi^+\right)^2+Y_{118}{}^{k_1\dot\alpha\beta}\ \overline\theta^2\,(D_\beta\Phi)\,\Phi\left(\partial_{k_1}\overline D_{\dot\alpha}\Phi^+\right)\left(\Phi^+\right)^2\\
&+Y_{119}{}^{k_1k_2}\ \overline\theta^2\ \ (\partial_{k_1}\partial_{k_2}\Phi)\,\Phi\left(\Phi^+\right)^3+Y_{120}{}^{k_1k_2}\ \overline\theta^2\ \ (\partial_{k_1}\Phi)\,(\partial_{k_2}\Phi)\left(\Phi^+\right)^3\\
&+Y_{121}{}^{k_1k_2}\ \overline\theta^2\ \Phi^2\left(\partial_{k_1}\partial_{k_2}\Phi^+\right)\left(\Phi^+\right)^2+Y_{122}\ \overline\theta^2\left(D^2\Phi\right)\left(D^2\Phi\right)\Phi\left(\overline D^2\Phi^+\right)\Phi^+\\
&+Y_{123}\ \epsilon^{\alpha\beta}\ \overline\theta^2\,(D_\alpha\Phi)\,(D_\beta\Phi)\left(D^2\Phi\right)\left(\overline D^2\Phi^+\right)\Phi^+\\
&+Y_{124}\ \epsilon^{\dot\alpha\dot\beta}\ \overline\theta^2\left(D^2\Phi\right)\left(D^2\Phi\right)\Phi\left(\overline D_{\dot\alpha}\Phi^+\right)\left(\overline D_{\dot\beta}\Phi^+\right)\\
&+Y_{125}{}^{k_1\dot\alpha\beta}\ \overline\theta^2\ \ (\partial_{k_1}D_\beta\Phi)\left(D^2\Phi\right)\Phi\left(\overline D_{\dot\alpha}\Phi^+\right)\Phi^++Y_{126}{}^{k_1\dot\alpha\beta}\ \overline\theta^2\,(D_\beta\Phi)\left(\partial_{k_1}D^2\Phi\right)\Phi\left(\overline D_{\dot\alpha}\Phi^+\right)\Phi^+\\
&+Y_{127}{}^{k_1\dot\alpha\beta}\ \overline\theta^2\,(D_\beta\Phi)\left(D^2\Phi\right)(\partial_{k_1}\Phi)\left(\overline D_{\dot\alpha}\Phi^+\right)\Phi^++Y_{128}{}^{k_1\dot\alpha\beta}\ \overline\theta^2\,(D_\beta\Phi)\left(D^2\Phi\right)\Phi\left(\partial_{k_1}\overline D_{\dot\alpha}\Phi^+\right)\Phi^+\\
&+Y_{129}{}^{k_1k_2}\ \overline\theta^2\ \ \left(\partial_{k_2}\partial_{k_1}D^2\Phi\right)\Phi^2\left(\Phi^+\right)^2+\overline Y_{130}{}^{k_1k_2}\ \overline\theta^2\ \ \left(\partial_{k_1}D^2\Phi\right)(\partial_{k_2}\Phi)\,\Phi\left(\Phi^+\right)^2\\
&+Y_{131}{}^{k_1k_2}\ \overline\theta^2\left(D^2\Phi\right)(\partial_{k_1}\partial_{k_2}\Phi)\,\Phi\left(\Phi^+\right)^2+Y_{132}{}^{k_1k_2}\ \overline\theta^2\left(D^2\Phi\right)(\partial_{k_1}\Phi)\,(\partial_{k_2}\Phi)\left(\Phi^+\right)^2\\
&+Y_{133}{}^{k_1k_2}\ \overline\theta^2\left(D^2\Phi\right)\Phi^2\left(\partial_{k_1}\partial_{k_2}\Phi^+\right)\Phi^++Y_{134}{}^{k_1k_2\alpha\beta}\ \overline\theta^2\ \ (\partial_{k_2}\partial_{k_1}D_\alpha\Phi)\,(D_\beta\Phi)\,\Phi\left(\Phi^+\right)^2\\
&+\overline Y_{135}{}^{k_1k_2\alpha\beta}\ \overline\theta^2\ \ (\partial_{k_1}D_\alpha\Phi)\,(\partial_{k_2}D_\beta\Phi)\,\Phi\left(\Phi^+\right)^2+\widetilde Y_{136}{}^{k_1k_2\alpha\beta}\ \overline\theta^2\ \ (\partial_{k_2}D_\beta\Phi)\,(D_\alpha\Phi)\,(\partial_{k_1}\Phi)\left(\Phi^+\right)^2\\
&+Y_{137}{}^{k_1k_2}\ \epsilon^{\alpha\beta}\ \overline\theta^2\,(D_\alpha\Phi)\,(D_\beta\Phi)\,(\partial_{k_1}\partial_{k_2}\Phi)\left(\Phi^+\right)^2\\
&+Y_{138}{}^{k_1k_2}\ \epsilon^{\alpha\beta}\ \overline\theta^2\,(D_\alpha\Phi)\,(D_\beta\Phi)\,\Phi\left(\partial_{k_1}\partial_{k_2}\Phi^+\right)\Phi^++Y_{139}\ \overline\theta^2\left(D^2\Phi\right)\left(D^2\Phi\right)\left(D^2\Phi\right)\Phi\left(\overline D^2\Phi^+\right)\\
&+Y_{140}\ \epsilon^{\alpha\beta}\ \overline\theta^2\,(D_\alpha\Phi)\,(D_\beta\Phi)\left(D^2\Phi\right)\left(D^2\Phi\right)\left(\overline D^2\Phi^+\right)\\
&+Y_{141}{}^{k_1\dot\alpha\beta}\ \overline\theta^2\ \ (\partial_{k_1}D_\beta\Phi)\left(D^2\Phi\right)\left(D^2\Phi\right)\Phi\left(\overline D_{\dot\alpha}\Phi^+\right)\\
&+Y_{142}{}^{k_1\dot\alpha\beta}\ \overline\theta^2\,(D_\beta\Phi)\left(\partial_{k_1}D^2\Phi\right)\left(D^2\Phi\right)\Phi\left(\overline D_{\dot\alpha}\Phi^+\right)\\
&+Y_{143}{}^{k_1\dot\alpha\beta}\ \overline\theta^2\,(D_\beta\Phi)\left(D^2\Phi\right)\left(D^2\Phi\right)(\partial_{k_1}\Phi)\left(\overline D_{\dot\alpha}\Phi^+\right)+Y_{144}{}^{k_1k_2}\ \overline\theta^2\ \ \left(\partial_{k_2}\partial_{k_1}D^2\Phi\right)\left(D^2\Phi\right)\Phi^2\ \Phi^+\\
&+Y_{145}{}^{k_1k_2}\ \overline\theta^2\ \ \left(\partial_{k_1}D^2\Phi\right)\left(\partial_{k_2}D^2\Phi\right)\Phi^2\ \Phi^++\overline Y_{146}{}^{k_1k_2}\ \overline\theta^2\ \ \left(\partial_{k_1}D^2\Phi\right)\left(D^2\Phi\right)(\partial_{k_2}\Phi)\,\Phi\ \Phi^+\\
&+Y_{147}{}^{k_1k_2}\ \overline\theta^2\left(D^2\Phi\right)\left(D^2\Phi\right)(\partial_{k_1}\partial_{k_2}\Phi)\,\Phi\ \Phi^++Y_{148}{}^{k_1k_2}\ \overline\theta^2\left(D^2\Phi\right)\left(D^2\Phi\right)(\partial_{k_1}\Phi)\,(\partial_{k_2}\Phi)\,\Phi^+\\
&+Y_{149}{}^{k_1k_2\alpha\beta}\ \overline\theta^2\ \ (\partial_{k_2}\partial_{k_1}D_\alpha\Phi)\,(D_\beta\Phi)\left(D^2\Phi\right)\Phi\ \Phi^+\\
&+\overline Y_{150}{}^{k_1k_2\alpha\beta}\ \overline\theta^2\ \ (\partial_{k_1}D_\alpha\Phi)\,(\partial_{k_2}D_\beta\Phi)\left(D^2\Phi\right)\Phi\ \Phi^+\\
&+\widetilde Y_{151}{}^{k_1k_2\alpha\beta}\ \overline\theta^2\ \ (\partial_{k_2}D_\beta\Phi)\,(D_\alpha\Phi)\left(D^2\Phi\right)(\partial_{k_1}\Phi)\,\Phi^+\\
&+\widetilde Y_{152}{}^{k_1k_2\alpha\beta}\ \overline\theta^2\ \ (\partial_{k_2}D_\beta\Phi)\,(D_\alpha\Phi)\left(\partial_{k_1}D^2\Phi\right)\Phi\ \Phi^+\\
&+Y_{153}{}^{k_1k_2}\ \epsilon^{\alpha\beta}\ \overline\theta^2\,(D_\alpha\Phi)\,(D_\beta\Phi)\left(\partial_{k_2}\partial_{k_1}D^2\Phi\right)\Phi\ \Phi^+
\end{aligned}
$$

$$
\begin{aligned}
&+\overline{Y}_{154}{}^{k_1k_2}\ \epsilon^{\alpha\beta}\ \overline{\theta}^2\ (D_\alpha\Phi)\ (D_\beta\Phi)\ \left(\partial_{k_1}D^2\Phi\right)\ (\partial_{k_2}\Phi)\ \Phi^+\\
&+Y_{155}{}^{k_1k_2}\ \epsilon^{\alpha\beta}\ \overline{\theta}^2\ (D_\alpha\Phi)\ (D_\beta\Phi)\ \left(D^2\Phi\right)\ (\partial_{k_1}\partial_{k_2}\Phi)\ \Phi^+ + Y_{156}\ \overline{\theta}^2\ \Phi\ \left(\overline{D}^2\Phi^+\right)\ \left(\Phi^+\right)^4\\
&+Y_{157}\ \epsilon^{\dot\alpha\dot\beta}\ \overline{\theta}^2\ \Phi\ \left(\overline{D}_{\dot\alpha}\Phi^+\right)\ \left(\overline{D}_{\dot\beta}\Phi^+\right)\ \left(\Phi^+\right)^3 + Y_{158}\ \overline{\theta}^2\ \left(D^2\Phi\right)\ \Phi\ \left(\overline{D}^2\Phi^+\right)\ \left(\Phi^+\right)^3\\
&+Y_{159}\ \epsilon^{\alpha\beta}\ \overline{\theta}^2\ (D_\alpha\Phi)\ (D_\beta\Phi)\ \left(\overline{D}^2\Phi^+\right)\ \left(\Phi^+\right)^3\\
&+Y_{160}\ \epsilon^{\dot\alpha\dot\beta}\ \overline{\theta}^2\ \left(D^2\Phi\right)\ \Phi\ \left(\overline{D}_{\dot\alpha}\Phi^+\right)\ \left(\overline{D}_{\dot\beta}\Phi^+\right)\ \left(\Phi^+\right)^2 + Y_{161}{}^{k_1\dot\alpha\dot\beta}\ \overline{\theta}^2\ \ (\partial_{k_1}D_\beta\Phi)\ \Phi\ \left(\overline{D}_{\dot\alpha}\Phi^+\right)\ \left(\Phi^+\right)^3\\
&+Y_{162}{}^{k_1\dot\alpha\beta}\ \overline{\theta}^2\ (D_\beta\Phi)\ (\partial_{k_1}\Phi)\ \left(\overline{D}_{\dot\alpha}\Phi^+\right)\ \left(\Phi^+\right)^3\\
&+Y_{163}{}^{k_1\dot\alpha\beta}\ \overline{\theta}^2\ (D_\beta\Phi)\ \Phi\ \left(\partial_{k_1}\overline{D}_{\dot\alpha}\Phi^+\right)\ \left(\Phi^+\right)^3 + Y_{164}{}^{k_1k_2}\ \overline{\theta}^2\ \ (\partial_{k_1}\partial_{k_2}\Phi)\ \Phi\ \left(\Phi^+\right)^4\\
&+Y_{165}{}^{k_1k_2}\ \overline{\theta}^2\ \ (\partial_{k_1}\Phi)\ (\partial_{k_2}\Phi)\ \left(\Phi^+\right)^4 + Y_{166}{}^{k_1k_2}\ \overline{\theta}^2\ \Phi^2\ \left(\partial_{k_1}\partial_{k_2}\Phi^+\right)\ \left(\Phi^+\right)^3\\
&+Y_{167}\ \overline{\theta}^2\ \left(D^2\Phi\right)\ \left(D^2\Phi\right)\ \Phi\ \left(\overline{D}^2\Phi^+\right)\ \left(\Phi^+\right)^2 + Y_{168}\ \epsilon^{\alpha\beta}\ \overline{\theta}^2\ (D_\alpha\Phi)\ (D_\beta\Phi)\ \left(D^2\Phi\right)\ \left(\overline{D}^2\Phi^+\right)\ \left(\Phi^+\right)^2\\
&+Y_{169}\ \epsilon^{\dot\alpha\dot\beta}\ \overline{\theta}^2\ \left(D^2\Phi\right)\ \left(D^2\Phi\right)\ \Phi\ \left(\overline{D}_{\dot\alpha}\Phi^+\right)\ \left(\overline{D}_{\dot\beta}\Phi^+\right)\ \Phi^+\\
&+Y_{170}{}^{k_1\dot\alpha\beta}\ \overline{\theta}^2\ \ (\partial_{k_1}D_\beta\Phi)\ \left(D^2\Phi\right)\ \Phi\ \left(\overline{D}_{\dot\alpha}\Phi^+\right)\ \left(\Phi^+\right)^2\\
&+Y_{171}{}^{k_1\dot\alpha\beta}\ \overline{\theta}^2\ (D_\beta\Phi)\ \left(\partial_{k_1}D^2\Phi\right)\ \Phi\ \left(\overline{D}_{\dot\alpha}\Phi^+\right)\ \left(\Phi^+\right)^2\\
&+Y_{172}{}^{k_1\dot\alpha\beta}\ \overline{\theta}^2\ (D_\beta\Phi)\ \left(D^2\Phi\right)\ (\partial_{k_1}\Phi)\ \left(\overline{D}_{\dot\alpha}\Phi^+\right)\ \left(\Phi^+\right)^2\\
&+Y_{173}{}^{k_1\dot\alpha\beta}\ \overline{\theta}^2\ (D_\beta\Phi)\ \left(D^2\Phi\right)\ \Phi\ \left(\partial_{k_1}\overline{D}_{\dot\alpha}\Phi^+\right)\ \left(\Phi^+\right)^2 + Y_{174}{}^{k_1k_2}\ \overline{\theta}^2\ \ \left(\partial_{k_2}\partial_{k_1}D^2\Phi\right)\ \Phi^2\ \left(\Phi^+\right)^3\\
&+\overline{Y}_{175}{}^{k_1k_2}\ \overline{\theta}^2\ \ \left(\partial_{k_2}D^2\Phi\right)\ (\partial_{k_1}\Phi)\ \Phi\ \left(\Phi^+\right)^3 + Y_{176}{}^{k_1k_2}\ \overline{\theta}^2\ \left(D^2\Phi\right)\ (\partial_{k_1}\partial_{k_2}\Phi)\ \Phi\ \left(\Phi^+\right)^3\\
&+Y_{177}{}^{k_1k_2}\ \overline{\theta}^2\ \left(D^2\Phi\right)\ (\partial_{k_1}\Phi)\ (\partial_{k_2}\Phi)\ \left(\Phi^+\right)^3 + Y_{178}{}^{k_1k_2}\ \overline{\theta}^2\ \left(D^2\Phi\right)\ \Phi\Phi\ \left(\partial_{k_1}\partial_{k_2}\Phi^+\right)\ \left(\Phi^+\right)^2\\
&+Y_{179}{}^{k_1k_2\alpha\beta}\ \overline{\theta}^2\ \ (\partial_{k_2}\partial_{k_1}D_\alpha\Phi)\ (D_\beta\Phi)\ \Phi\ \left(\Phi^+\right)^3\\
&+\overline{Y}_{180}{}^{k_1k_2\alpha\beta}\ \overline{\theta}^2\ \ (\partial_{k_1}D_\alpha\Phi)\ (\partial_{k_2}D_\beta\Phi)\ \Phi\ \left(\Phi^+\right)^3 + \widetilde{Y}_{181}{}^{k_1k_2\alpha\beta}\ \overline{\theta}^2\ \ (\partial_{k_2}D_\beta\Phi)\ (D_\alpha\Phi)\ (\partial_{k_1}\Phi)\ \left(\Phi^+\right)^3\\
&+Y_{182}{}^{k_1k_2}\ \epsilon^{\alpha\beta}\ \overline{\theta}^2\ (D_\alpha\Phi)\ (D_\beta\Phi)\ (\partial_{k_1}\partial_{k_2}\Phi)\ \left(\Phi^+\right)^3\\
&+Y_{183}{}^{k_1k_2}\ \epsilon^{\alpha\beta}\ \overline{\theta}^2\ (D_\alpha\Phi)\ (D_\beta\Phi)\ \Phi\ \left(\partial_{k_1}\partial_{k_2}\Phi^+\right)\ \left(\Phi^+\right)^2\\
&+Y_{184}\ \overline{\theta}^2\ \Phi\ \left(\overline{D}^2\Phi^+\right)\ \left(\Phi^+\right)^5 + Y_{185}\ \epsilon^{\dot\alpha\dot\beta}\ \overline{\theta}^2\ \Phi\ \left(\overline{D}_{\dot\alpha}\Phi^+\right)\ \left(\overline{D}_{\dot\beta}\Phi^+\right)\ \left(\Phi^+\right)^4\\
&+Y_{186}\ \overline{\theta}^2\ \left(D^2\Phi\right)\ \Phi\ \left(\overline{D}^2\Phi^+\right)\ \left(\Phi^+\right)^4 + Y_{187}\ \epsilon^{\alpha\beta}\ \overline{\theta}^2\ (D_\alpha\Phi)\ (D_\beta\Phi)\ \left(\overline{D}^2\Phi^+\right)\ \left(\Phi^+\right)^4\\
&+Y_{188}\ \epsilon^{\dot\alpha\dot\beta}\ \overline{\theta}^2\ \left(D^2\Phi\right)\ \Phi\ \left(\overline{D}_{\dot\alpha}\Phi^+\right)\ \left(\overline{D}_{\dot\beta}\Phi^+\right)\ \left(\Phi^+\right)^3 + Y_{189}{}^{k_1\dot\alpha\beta}\ \overline{\theta}^2\ \ (\partial_{k_1}D_\beta\Phi)\ \Phi\ \left(\overline{D}_{\dot\alpha}\Phi^+\right)\ \left(\Phi^+\right)^4\\
&+Y_{190}{}^{k_1\dot\alpha\beta}\ \overline{\theta}^2\ (D_\beta\Phi)\ (\partial_{k_1}\Phi)\ \left(\overline{D}_{\dot\alpha}\Phi^+\right)\ \left(\Phi^+\right)^4 + Y_{191}{}^{k_1\dot\alpha\beta}\ \overline{\theta}^2\ (D_\beta\Phi)\ \Phi\ \left(\partial_{k_1}\overline{D}_{\dot\alpha}\Phi^+\right)\ \left(\Phi^+\right)^4\\
&+Y_{192}{}^{k_1k_2}\ \overline{\theta}^2\ \ (\partial_{k_1}\partial_{k_2}\Phi)\ \Phi\ \left(\Phi^+\right)^5 + Y_{193}{}^{k_1k_2}\ \overline{\theta}^2\ \ (\partial_{k_1}\Phi)\ (\partial_{k_2}\Phi)\ \left(\Phi^+\right)^5\\
&+Y_{194}{}^{k_1k_2}\ \overline{\theta}^2\ \Phi^2\ \left(\partial_{k_1}\partial_{k_2}\Phi^+\right)\ \left(\Phi^+\right)^4 + Y_{195}\ \overline{\theta}^2\ \Phi\ \left(\overline{D}^2\Phi^+\right)\ \left(\Phi^+\right)^6\\
&+Y_{196}\ \epsilon^{\dot\alpha\dot\beta}\ \overline{\theta}^2\ \Phi\ \left(\overline{D}_{\dot\alpha}\Phi^+\right)\ \left(\overline{D}_{\dot\beta}\Phi^+\right)\ \left(\Phi^+\right)^5\ \Bigg\}. \qquad \text{(A38)}
\end{aligned}
$$

References

[1] M. R. Douglas and N. A. Nekrasov, "Noncommutative field theory," Rev. Mod. Phys. **73**, 977 (2001) [arXiv:hep-th/0106048].

[2] R. J. Szabo, "Quantum field theory on noncommutative spaces," Phys. Rept. **378**, 207 (2003) [arXiv:hep-th/0109162].

[3] N. Seiberg and E. Witten, "String theory and noncommutative geometry," JHEP **9909**, 032 (1999) [arXiv:hep-th/9908142].

[4] S. Ferrara, M. A. Lledo and O. Macia, "Supersymmetry in noncommutative superspaces," JHEP **0309**, 068 (2003) [hep-th/0307039].

[5] H. O. Girotti, M. Gomes, V. O. Rivelles and A. J. da Silva, "A Consistent noncommutative field theory: The Wess-Zumino model," Nucl. Phys. B **587**, 299 (2000) [hep-th/0005272].

[6] H. Ooguri and C. Vafa, "The C-deformation of gluino and nonplanar diagrams," Adv. Theor. Math. Phys. **7**, 53 (2003) [arXiv:hep-th/0302109].

[7] H. Ooguri and C. Vafa, "Gravity induced C-deformation," Adv. Theor. Math. Phys. **7**, 405 (2004) [arXiv:hep-th/0303063].

[8] N. Seiberg, "Noncommutative superspace, N = 1/2 supersymmetry, field theory and string theory," JHEP **0306**, 010 (2003) [arXiv:hep-th/0305248].

[9] M. T. Grisaru, S. Penati, and A. Romagnoni, "Two loop renormalization for nonanticommutative N = 1/2 supersymmetric WZ model," JHEP **0308**, 003 (2003) [arXiv:hep-th/0307099].

[10] R. Britto and B. Feng, "N=1/2 Wess-Zumino model is renormalizable," Phys. Rev. Lett. **91**, 201601 (2003) [arXiv:hep-th/0307165].

[11] A. Romagnoni, "Renormalizability of N=1/2 Wess-Zumino model in superspace," JHEP **0310**, 016 (2003) [arXiv:hep-th/0307209].

[12] J. de Boer, P. A. Grassi, and P. van Nieuwenhuizen, "Noncommutative superspace from string theory," Phys. Lett. B **574**, 98 (2003) [arXiv:hep-th/0302078].

[13] Y. G. Miao and X. D. Wang, "One-loop renormalizable Wess-Zumino model on a bosonic-fermionic noncommutative superspace," Phys. Rev. D **90**, no. 4, 045036 (2014) [arXiv:1403.4705 [hep-th]].

[14] Y. G. Miao and X. D. Wang, "All-Loop Renormalizable Wess-Zumino Model on Bosonic-Fermionic Noncommutative Superspace," Phys. Rev. D **91**, no. 2, 025016 (2015) [arXiv:1403.5046 [hep-th]].

[15] M. T. Grisaru, S. Penati and A. Romagnoni, "Non(anti)commutative sym theory: Renormalization in superspace," JHEP **0602**, 043 (2006) [hep-th/0510175].

[16] X. D. Wang, "Covariant Quantization of BFNC Super Yang-Mills Theories and Supergauge Invariance ," [arXiv:1603.04526].